

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
25 March 2004 (25.03.2004)

PCT

(10) International Publication Number  
**WO 2004/024700 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 239/22**,  
401/04, 401/12, 403/08, 403/12, 405/08, 413/04, 417/04,  
A61K 31/513, 31/5377, A61P 9/00

(21) International Application Number:  
PCT/EP2003/009525

(22) International Filing Date: 28 August 2003 (28.08.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0220962.5 10 September 2002 (10.09.2002) GB  
0226609.6 14 November 2002 (14.11.2002) GB  
0315870.6 7 July 2003 (07.07.2003) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): BAYER HEALTHCARE AG [DE/DE]; 51368 Leverkusen (DE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): GIELEN, Heike [DE/DE]; Am Kettnersbusch 3, 51379 Leverkusen (DE). LI, Volkhart [DE/DE]; Im Wiesengrund 40, 42553 Velbert (DE). ROSENTER, Ulrich [DE/DE]; Obere Rutenbeck 6, 42349 Wuppertal (DE). SCHLEMMER, Karl-Heinz [DE/DE]; Wildsteig 22a, 42113 Wuppertal (DE). ALLERHEILIGEN, Swen [DE/DE]; Beckumsfeld 4, 45259 Essen (DE). TELAN, Leila [US/DE]; Rabenweg 42, 42115 Wuppertal (DE). BÄRFACKER, Lars [DE/DE]; Bachstr. 98, 46149 Oberhausen (DE). KELDENICH, Jörg [DE/DE]; Damaschkeweg 49, 42113 Wuppertal (DE). FITZGERALD, Mary, F. [GB/GB]; 2 Paternoster Court, Cassington Road, Yarnton, Oxfordshire OX5 1QB (GB). NASH, Kevin [GB/GB]; 136A Littlebrook Avenue, Slough, Burnham, Berkshire SL2 2NE (GB). ALBRECHT, Barbara [DE/DE]; Heidestrasse 9, 42489 Wülfrath (DE). MEURER, Dirk [DE/DE]; Platanenweg 1a, 50259 Pulheim (DE).

(74) Common Representative: BAYER HEALTHCARE AG; Law and Patents, Patents and Licensing, 51368 Leverkusen (DE).

#### Declaration under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

#### Published:

— with international search report  
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PYRIMIDINONE DERIVATIVES AS THERAPEUTIC AGENTS AGAINST ACUTE AND CHRONIC INFLAMMATORY, ISCHAEMIC AND REMODELLING PROCESSES

(57) Abstract: The invention relates to novel heterocyclic derivatives, processes for their preparation, and their use in medicaments, especially for the treatment of chronic obstructive pulmonary diseases.

WO 2004/024700 A1

- 1 -

## PYRIMIDINONE DERIVATIVES AS THERAPEUTIC AGENTS AGAINST ACUTE AND CHRONIC INFLAMMATORY, ISCHAEMIC AND REMODELLING PROCESSES

The present invention relates to novel heterocyclic derivatives, processes for their preparation, and their use in medicaments, especially for the treatment of chronic  
5 obstructive pulmonary diseases, acute coronary syndrome, acute myocardial infarction and heart failure development.

The fibrous protein elastin, which comprises an appreciable percentage of all protein content in some tissues, such as the arteries, some ligaments, the lungs and the heart,  
10 can be hydrolysed or otherwise destroyed by a select group of enzymes classified as elastases. Human leukocyte elastase (HLE, EC 3.4.21.37), also known as human neutrophil elastase (HNE), is a glycosylated, strongly basic serine protease and is found in the azurophilic granules of human polymorphonuclear leukocytes (PMN). HNE is released from activated PMN and has been implicated causally in the  
15 pathogenesis of acute and chronic inflammatory diseases. HNE is capable of degrading a wide range of matrix proteins including elastin and collagen, and in addition to these actions on connective tissue HNE has a broad range of inflammatory actions including upregulation of IL-8 gene expression, oedema formation, mucus gland hyperplasia and mucus hypersecretion. It also acts as a  
20 mediator of tissue injury by hydrolysing collagen structures, e.g. in the heart after acute myocardial infarction or during the development of heart failure, thus damaging endothelial cells, promoting extravasation of neutrophils adhering to the endothelium and influencing the adhesion process itself.

25 Pulmonary diseases where HNE is believed to play a role include lung fibrosis, pneumonia, acute respiratory distress syndrome (ARDS), pulmonary emphysema, including smoking-induced emphysema, chronic obstructive pulmonary diseases (COPD) and cystic fibrosis. In cardiovascular diseases, HNE is involved in the enhanced generation of ischaemic tissue injury followed by myocardial dysfunction  
30 after acute myocardial infarction and in the remodelling processes occurring during the development of heart failure. HNE has also been causally implicated in

- 2 -


rheumatoid arthritis, atherosclerosis, brain trauma, cancer and related conditions in which neutrophil participation is involved.

Thus, inhibitors of HLE activity can be potentially useful in the treatment of a number of inflammatory diseases, especially of chronic obstructive pulmonary diseases [R.A. Stockley, *Neutrophils and protease/antiprotease imbalance*, Am. J. Respir. Crit. Care 160, S49-S52 (1999)]. Inhibitors of HLE activity can also be potentially useful in the treatment of acute myocardial syndrome, unstable angina pectoris, acute myocardial infarction and coronary artery bypass grafts (CABG) [C.P. Tiefenbacher et al., *Inhibition of elastase improves myocardial function after repetitive ischaemia and myocardial infarction in the rat heart*, Eur. J. Physiol. 433, S563-S570 (1997); Dinerman et al., *Increased neutrophil elastase release in unstable angina pectoris and acute myocardial infarction*, J. Am. Coll. Cardiol. 15, 1559-1563 (1990)], of the development of heart failure [S.J. Gilbert et al., *Increased expression of promatrix metalloproteinase-9 and neutrophil elastase in canine dilated cardiomyopathy*, Cardiovasc. Res. 34, S377-S383 (1997)] and of atherosclerosis [Dollery et al., *Neutrophil elastase in human atherosclerotic plaque*, Circulation 107, 2829-2836 (2003)].

The synthesis of 5-ethoxycarbonyl-1-phenyl-6-methyl-4-(3-nitrophenyl)-3,4-dihydropyrimidin-2(1H)-one is described in J. Heterocyclic Chem. 38, 1051 (2001). A pharmacological activity of this compound is not mentioned.

The present invention relates to compounds of the general formula (I)

25



wherein

A represents an aryl or heteroaryl ring,

5

$R^1$ ,  $R^2$  and  $R^3$  independently from each other represent hydrogen, halogen, nitro, cyano,  $C_1$ - $C_6$ -alkyl, hydroxy or  $C_1$ - $C_6$ -alkoxy, wherein  $C_1$ - $C_6$ -alkyl and  $C_1$ - $C_6$ -alkoxy can be further substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy and  $C_1$ - $C_4$ -alkoxy,

10

$R^4$  represents trifluoromethylcarbonyl,  $C_1$ - $C_6$ -alkylcarbonyl,  $C_1$ - $C_6$ -alkoxy-carbonyl,  $C_1$ - $C_6$ -alkenoxycarbonyl, hydroxycarbonyl, aminocarbonyl, mono- or di- $C_1$ - $C_4$ -alkylaminocarbonyl,  $C_6$ - $C_{10}$ -arylaminocarbonyl, arylcarbonyl, heteroarylcarbonyl, heterocyclylcarbonyl, heteroaryl, heterocyclyl or cyano, wherein  $C_1$ - $C_6$ -alkylcarbonyl,  $C_1$ - $C_6$ -alkoxycarbonyl, mono- and di- $C_1$ - $C_4$ -alkylaminocarbonyl can be further substituted with one to three identical or different radicals selected from the group consisting of  $C_3$ - $C_8$ -cycloalkyl, hydroxy,  $C_1$ - $C_4$ -alkoxy,  $C_1$ - $C_4$ -alkoxycarbonyl, hydroxycarbonyl, amino-carbonyl, mono- and di- $C_1$ - $C_4$ -alkylaminocarbonyl,  $C_1$ - $C_4$ -alkylcarbonyl-amino, ( $C_1$ - $C_4$ -alkylcarbonyl)- $C_1$ - $C_4$ -alkylamino, cyano, amino, mono- and di- $C_1$ - $C_4$ -alkylamino, heteroaryl, heterocyclyl and tri-( $C_1$ - $C_6$ -alkyl)-silyl, and

20



wherein heteroarylcarbonyl, heterocyclylcarbonyl, heteroaryl and heterocyclyl can be further substituted with C<sub>1</sub>-C<sub>4</sub>-alkyl,

5           R<sup>5</sup>       represents C<sub>1</sub>-C<sub>4</sub>-alkyl, which can be substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-alkenoxy, C<sub>1</sub>-C<sub>6</sub>-alkylthio, amino, mono- and di-C<sub>1</sub>-C<sub>6</sub>-alkylamino, arylamino, hydroxycarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl and the radical -O-C<sub>1</sub>-C<sub>4</sub>-alkyl-O-C<sub>1</sub>-C<sub>4</sub>-alkyl,

10       or


          R<sup>5</sup>       represents amino,

15       R<sup>6</sup>       represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, formyl, aminocarbonyl, mono- or di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, N-(C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl)-aminocarbonyl, N-(C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl)-N-(C<sub>1</sub>-C<sub>4</sub>-alkyl)-aminocarbonyl, heteroaryl, heterocyclyl, heteroarylcarbonyl or heterocyclylcarbonyl, wherein C<sub>1</sub>-C<sub>6</sub>-alkyl, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, 20       heteroaryl and heterocyclyl can be substituted with one to three identical or different radicals selected from the group consisting of aryl, heteroaryl, hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, hydroxycarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, aminocarbonyl, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, amino, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>1</sub>-C<sub>4</sub>-alkylcarbonylamino, tri-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-silyl, cyano, 25       N-(mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylamino-C<sub>1</sub>-C<sub>4</sub>-alkyl)-aminocarbonyl, N-(C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl)-aminocarbonyl and halogen,

          or

30       R<sup>6</sup>       represents a moiety of the formula

- 5 -



wherein

$R^{6A}$  is selected from the group consisting of hydrogen and  $C_1$ - $C_6$ -alkyl, and

$n$  represents an integer of 1 or 2,

$R^7$  represents halogen, nitro, cyano,  $C_1$ - $C_6$ -alkyl, hydroxy or  $C_1$ - $C_6$ -alkoxy, wherein  $C_1$ - $C_6$ -alkyl and  $C_1$ - $C_6$ -alkoxy can be further substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy and  $C_1$ - $C_4$ -alkoxy,

and

$Y^1$ ,  $Y^2$ ,  $Y^3$ ,  $Y^4$  and  $Y^5$  independently from each other represent CH or N, wherein the ring contains either 0, 1 or 2 nitrogen atoms.

The compounds according to this invention can also be present in the form of their salts, hydrates and/or solvates.

Physiologically acceptable salts are preferred in the context of the present invention.

Physiologically acceptable salts according to the invention are non-toxic salts which in general are accessible by reaction of the compounds (I) with an inorganic or organic base or acid conventionally used for this purpose. Non-limiting examples of pharmaceutically acceptable salts of compounds (I) include the alkali metal salts, e.g. lithium, potassium and sodium salts, the alkaline earth metal salts such as magnesium and calcium salts, the quaternary ammonium salts such as, for example, triethyl ammonium salts, acetates, benzene sulphonates, benzoates, dicarbonates, disulphates,

5     ditartrates, borates, bromides, carbonates, chlorides, citrates, dihydrochlorides, fumarates, gluconates, glutamates, hexyl resorcinates, hydrobromides, hydrochlorides, hydroxynaphthoates, iodides, isothionates, lactates, laurates, malates, maleates, mandelates, mesylates, methylbromides, methylnitrates, methylsulphates, nitrates, oleates, oxalates, palmitates, pantothenates, phosphates, diphosphates, polygalacturonates, salicylates, stearates, sulphates, succinates, tartrates, tosylates, valerates, and other salts used for medicinal purposes.

10     Hydrates of the compounds of the invention or their salts are stoichiometric compositions of the compounds with water, such as for example hemi-, mono-, or dihydrates.

15     Solvates of the compounds of the invention or their salts are stoichiometric compositions of the compounds with solvents.

20     The present invention includes both the individual enantiomers or diastereomers and the corresponding racemates or diastereomeric mixtures of the compounds according to the invention and their respective salts. In addition, all possible tautomeric forms of the compounds described above are included according to the present invention.

25     The diastereomeric mixtures can be separated into the individual isomers by chromatographic processes. The racemates can be resolved into the respective enantiomers either by chromatographic processes on chiral phases or by resolution.

30     In the context of the present invention, the substituents, if not stated otherwise, in general have the following meaning:

35     Alkyl in general represents a straight-chain or branched hydrocarbon radical having 1 to 6, preferably 1 to 4 carbon atoms. Non-limiting examples include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec.-butyl, tert.-butyl, pentyl, isopentyl, hexyl, isohexyl. The same applies to radicals such as alkoxy, alkylamino, alkoxycarbonyl and alkoxycarbonylamino.

Alkoxy illustratively and preferably represents methoxy, ethoxy, n-propoxy, isopropoxy, tert.-butoxy, n-pentoxy and n-hexoxy.

5     Alkylcarbonyl in general represents a straight-chain or branched hydrocarbon radical having 1 to 6, preferably 1 to 4 carbon atoms which has a carbonyl function at the position of attachment. Non-limiting examples include formyl, acetyl, n-propionyl, n-butyryl, isobutyryl, pivaloyl, n-hexanoyl.

10    Alkoxy carbonyl illustratively and preferably represents methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl, isopropoxycarbonyl, tert.-butoxycarbonyl, n-pentoxycarbonyl and n-hexoxycarbonyl.

15    Alkylamino represents an alkylamino radical having one or two (independently selected) alkyl substituents, illustratively and preferably representing methylamino, ethylamino, n-propylamino, isopropylamino, tert.-butylamino, n-pentylamino, n-hexylamino, *N,N*-dimethylamino, *N,N*-diethylamino, *N*-ethyl-*N*-methylamino, *N*-methyl-*N*-n-propylamino, *N*-isopropyl-*N*-n-propylamino, *N*-tert.-butyl-*N*-methylamino, *N*-ethyl-*N*-n-pentylamino and *N*-n-hexyl-*N*-methylamino.

20    Alkylaminocarbonyl represents an alkylaminocarbonyl radical having one or two (independently selected) alkyl substituents, illustratively and preferably representing methylaminocarbonyl, ethylaminocarbonyl, n-propylaminocarbonyl, isopropylaminocarbonyl, tert.-butylaminocarbonyl, n-pentylaminocarbonyl, n-hexylaminocarbonyl, *N,N*-dimethylaminocarbonyl, *N,N*-diethylaminocarbonyl, *N*-ethyl-*N*-methylaminocarbonyl, *N*-methyl-*N*-n-propylaminocarbonyl, *N*-isopropyl-*N*-n-propylaminocarbonyl, *N*-tert.-butyl-*N*-methylaminocarbonyl, *N*-ethyl-*N*-n-pentylaminocarbonyl and *N*-n-hexyl-*N*-methylaminocarbonyl.

30    Alkylsulfonyl in general represents a straight-chain or branched hydrocarbon radical having 1 to 6, preferably 1 to 4 carbon atoms which has a sulfonyl function at the

position of attachment. Non-limiting examples include methylsulfonyl, ethylsulfonyl, n-propylsulfonyl, isopropylsulfonyl, n-butylsulfonyl, tert.-butylsulfonyl.

5     Cycloalkyl in general represents a cyclic saturated hydrocarbon radical having 3 to 8, preferably 3 to 6 carbon atoms. Non-limiting examples include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

10     Aryl per se and in arylcarbonyl represents a mono- to tricyclic aromatic carbocyclic radical having generally 6 to 14 carbon atoms, illustratively and preferably representing phenyl, naphthyl and phenanthrenyl.

Arylcarbonyl illustratively and preferably represents benzoyl and naphthoyl.

15     Heteroaryl per se and in heteroarylcarbonyl represents an aromatic mono- or bicyclic radical having generally 5 to 10 and preferably 5 or 6 ring atoms and up to 5 and preferably up to 4 heteroatoms selected from the group consisting of S, O and N, illustratively and preferably representing thienyl, furyl, pyrrolyl, thiazolyl, oxazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, pyridyl, pyrimidyl, pyridazinyl, indolyl, indazolyl, benzofuranyl, benzothiophenyl, benzothiazolyl, quinolinyl, isoquinolinyl.

20     Heteroarylcarbonyl illustratively and preferably represents thienylcarbonyl, furylcarbonyl, pyrrolylcarbonyl, thiazolylcarbonyl, oxazolylcarbonyl, imidazolylcarbonyl, pyridylcarbonyl, pyrimidylcarbonyl, pyridazinylcarbonyl, indolylcarbonyl, indazolylcarbonyl, benzofuranylcarbonyl, benzothiophenylcarbonyl, quinolinylcarbonyl, isoquinolinylcarbonyl.

25

30     Heterocyclyl per se and in heterocyclylcarbonyl represents a mono- or polycyclic, preferably mono- or bicyclic, nonaromatic heterocyclic radical having generally 4 to 10 and preferably 5 to 8 ring atoms and up to 3 and preferably up to 2 heteroatoms and/or hetero groups selected from the group consisting of N, O, S, SO and SO<sub>2</sub>. The heterocyclyl radicals can be saturated or partially unsaturated. Preference is given to

5- to 8-membered monocyclic saturated heterocyclyl radicals having up to two heteroatoms selected from the group consisting of O, N and S, such as illustratively and preferably tetrahydrofuran-2-yl, pyrrolidin-1-yl, pyrrolidin-2-yl, pyrrolidin-3-yl, pyrrolinyl, piperidinyl, morpholinyl, perhydroazepinyl.

5

Heterocyclylcarbonyl illustratively and preferably represents tetrahydrofuran-2-carbonyl, pyrrolidine-1-carbonyl, pyrrolidine-2-carbonyl, pyrrolidine-3-carbonyl, pyrrolinecarbonyl, piperidinecarbonyl, morpholinecarbonyl, perhydroazepinecarbonyl.

10

Halogen represents fluorine, chlorine, bromine and iodine.

When stated, that  $Y^1$ ,  $Y^2$ ,  $Y^3$ ,  $Y^4$  and  $Y^5$  represent CH or N, CH shall also stand for a ring carbon atom, which is substituted with a substituent  $R^3$  or  $R^7$ .

15

A \* symbol next to a bond denotes the point of attachment in the molecule.

In another embodiment, the present invention relates to compounds of general formula (I), wherein

20

A represents an aryl or heteroaryl ring,

$R^1$ ,  $R^2$  and  $R^3$  independently from each other represent hydrogen, halogen, nitro, cyano,  $C_1$ - $C_6$ -alkyl, hydroxy or  $C_1$ - $C_6$ -alkoxy, wherein  $C_1$ - $C_6$ -alkyl and  $C_1$ - $C_6$ -alkoxy can be further substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy and  $C_1$ - $C_4$ -alkoxy,

25

$R^4$  represents  $C_1$ - $C_6$ -alkylcarbonyl,  $C_1$ - $C_6$ -alkoxycarbonyl,  $C_1$ - $C_6$ -alkenoxycarbonyl, hydroxycarbonyl, aminocarbonyl, mono- or di- $C_1$ - $C_4$ -alkylamino-carbonyl,  $C_6$ - $C_{10}$ -arylaminocarbonyl, heteroarylcarbonyl, heterocyclyl-

30

carbonyl, heteroaryl, heterocyclyl or cyano, wherein C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl can be further substituted with one to three identical or different radicals selected from the group consisting of C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl, hydroxycarbonyl, aminocarbonyl, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, C<sub>1</sub>-C<sub>4</sub>-alkylcarbonylamino, amino, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylamino, heteroaryl, heterocyclyl and tri-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-silyl,

R<sup>5</sup> represents C<sub>1</sub>-C<sub>4</sub>-alkyl, which can be substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-alkenoxy, C<sub>1</sub>-C<sub>6</sub>-alkylthio, amino, mono- and di-C<sub>1</sub>-C<sub>6</sub>-alkylamino, arylamino, hydroxycarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl and the radical -O-C<sub>1</sub>-C<sub>4</sub>-alkyl-O-C<sub>1</sub>-C<sub>4</sub>-alkyl,

or


R<sup>5</sup> represents amino,

R<sup>6</sup> represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, formyl, aminocarbonyl, mono- or di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, N-(C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl)-aminocarbonyl, N-(C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl)-N-(C<sub>1</sub>-C<sub>4</sub>-alkyl)-aminocarbonyl, heteroaryl, heterocyclyl, heteroarylcarbonyl or heterocyclylcarbonyl, wherein C<sub>1</sub>-C<sub>6</sub>-alkyl, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, heteroaryl and heterocyclyl can be substituted with one to three identical or different radicals selected from the group consisting of aryl, heteroaryl, hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, hydroxycarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, aminocarbonyl, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, amino, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylamino, C<sub>1</sub>-C<sub>4</sub>-alkylcarbonylamino, tri-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-silyl, cyano, N-(mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylamino-C<sub>1</sub>-C<sub>4</sub>-alkyl)-aminocarbonyl, N-(C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkyl)-aminocarbonyl and halogen,

or

$R^6$  represents a moiety of the formula

5



wherein

$R^{6A}$  is selected from the group consisting of hydrogen and  $C_1$ - $C_6$ -alkyl, and

10

$n$  represents an integer of 1 or 2,

$R^7$  represents halogen, nitro, cyano,  $C_1$ - $C_6$ -alkyl, hydroxy or  $C_1$ - $C_6$ -alkoxy, wherein  $C_1$ - $C_6$ -alkyl and  $C_1$ - $C_6$ -alkoxy can be further substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy and  $C_1$ - $C_4$ -alkoxy,

15

and

$Y^1$ ,  $Y^2$ ,  $Y^3$ ,  $Y^4$  and  $Y^5$  independently from each other represent CH or N, wherein the ring contains either 0, 1 or 2 nitrogen atoms.

20

In another embodiment, the present invention relates to compounds of general formula (I), wherein

25

A represents a phenyl, naphthyl or pyridyl ring,

$R^1$ ,  $R^2$  and  $R^3$  independently from each other represent hydrogen, fluoro, chloro, bromo, nitro, cyano, methyl, ethyl, trifluoromethyl or trifluoromethoxy,




R<sup>4</sup> represents C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, hydroxycarbonyl, aminocarbonyl, mono-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl or cyano, wherein C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl and mono-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl  
 5 can be substituted with one to three identical or different radicals selected from the group consisting of C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl, amino, mono- or di-C<sub>1</sub>-C<sub>4</sub>-alkylamino, heteroaryl and heterocyclyl,

10 R<sup>5</sup> represents methyl or ethyl,

R<sup>6</sup> represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, mono- or di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl or heterocyclcarbonyl, wherein C<sub>1</sub>-C<sub>6</sub>-alkyl and C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl can be substituted with one to three  
 15 identical or different radicals selected from the group consisting of heteroaryl, hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, hydroxycarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, aminocarbonyl, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, cyano, amino, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylamino,

20 or

R<sup>6</sup> represents a moiety of the formula



25 wherein

R<sup>6A</sup> is selected from the group consisting of hydrogen and C<sub>1</sub>-C<sub>4</sub>-alkyl, and

n represents an integer of 1 or 2,

R<sup>7</sup> represents halogen, nitro, cyano, trifluoromethyl, trifluoromethoxy, methyl or ethyl,

5 and

Y<sup>1</sup>, Y<sup>2</sup>, Y<sup>3</sup>, Y<sup>4</sup> and Y<sup>5</sup> each represent CH.

10 In another embodiment, the present invention relates to compounds of general formula (I), wherein

A represents a phenyl or a pyridyl ring,

15 R<sup>1</sup> and R<sup>3</sup> each represent hydrogen,

R<sup>2</sup> represents fluoro, chloro, bromo, nitro or cyano,


20 R<sup>4</sup> represents cyano, C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyl or C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl, wherein C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl can be substituted with a radical selected from the group consisting of hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylamino, heteroaryl and heterocyclyl,

R<sup>5</sup> represents methyl,

25 R<sup>6</sup> represents hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, mono- or di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyl or C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl, wherein C<sub>1</sub>-C<sub>4</sub>-alkyl and C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl can be substituted with a radical selected from the group consisting of heteroaryl, hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, hydroxycarbonyl, aminocarbonyl, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, amino, mono- and  
30 di-C<sub>1</sub>-C<sub>4</sub>-alkylamino,

or

$R^6$  represents a moiety of the formula



wherein

$R^{6A}$  is selected from the group consisting of hydrogen and methyl,

10  $R^7$  represents trifluoromethyl or nitro,

and

$Y^1, Y^2, Y^3, Y^4$  and  $Y^5$  each represent CH.

15

In another embodiment, the present invention relates to compounds according to general formula (I), wherein A is phenyl or pyridyl.

20 In another embodiment, the present invention relates to compounds according to general formula (I), wherein  $R^1$  is hydrogen.

25 In another embodiment, the present invention relates to compounds according to general formula (I), wherein  $R^2$  is cyano, especially wherein A is phenyl or pyridyl and  $R^2$  is cyano located in para-position relative to the central dihydropyrimidinone ring.

In another embodiment, the present invention relates to compounds according to general formula (I), wherein  $R^3$  is hydrogen.

In another embodiment, the present invention relates to compounds according to general formula (I), wherein  $R^4$  is  $C_1$ - $C_4$ -alkoxycarbonyl optionally substituted by hydroxy, especially 2-hydroxyethoxycarbonyl, or wherein  $R^4$  is  $C_1$ - $C_4$ -alkylcarbonyl, especially methylcarbonyl.

5

In another embodiment, the present invention relates to compounds according to general formula (I), wherein  $R^5$  is methyl.


10

In another embodiment, the present invention relates to compounds according to general formula (I), wherein  $R^6$  is hydrogen.

15

In another embodiment, the present invention relates to compounds according to general formula (I), wherein  $R^7$  is trifluoromethyl or nitro, especially wherein  $R^7$  is trifluoromethyl located in meta-position relative to the central dihydropyrimidinone ring.

In another embodiment, the present invention relates to compounds of general formula (IA)



20


wherein

$Z$  represents CH or N, and

$R^1$ ,  $R^3$ ,  $R^4$  and  $R^6$  have the meaning indicated above.


The compounds of the present invention, wherein  $R^6$  is hydrogen, can enolize into the corresponding hydroxyamidines:

5



The compounds of general formula (I) can be synthesized by condensing compounds of general formula (II)

10




wherein

$A$ ,  $R^1$  and  $R^2$  have the meaning indicated above,

15

with compounds of general formula (III)

- 17 -




wherein

R<sup>4</sup> and R<sup>5</sup> have the meaning indicated above,

5

and compounds of general formula (IV)




wherein

10

R<sup>3</sup>, R<sup>7</sup>, and Y<sup>1</sup> to Y<sup>5</sup> have the meaning indicated above,

in the presence of an acid either in a three-component / one-step reaction or sequentially to give compounds of the general formula (IB)

15



wherein

A, R<sup>1</sup> to R<sup>5</sup>, R<sup>7</sup>, and Y<sup>1</sup> to Y<sup>5</sup> have the meaning indicated above,

optionally followed by reaction of the compounds of general formula (IB) with  
5 compounds of the general formula (V)



wherein

10

R<sup>6\*</sup> has the meaning of R<sup>6</sup> as indicated above, but does not represent hydrogen,  
and

15

X represents a leaving group, such as halogen, tosylate, mesylate or sulfate,

in the presence of a base.

20

The compounds of general formula (I), wherein R<sup>4</sup> represents cyano, R<sup>5</sup> represents amino and R<sup>6</sup> represents hydrogen, can alternatively be prepared by condensing  
compounds of general formula (II) with compounds of general formula (IV) and a  
compound of formula (VI)



25

in the presence of an acid either in a three-component / one-step reaction or sequentially.

30

Suitable solvents for the process (II) + (III)/(VI) + (IV) → (IB) are generally customary organic solvents which do not change under the reaction conditions. These include ethers such as diethyl ether, diisopropyl ether, 1,2-dimethoxyethane, dioxan or tetrahydrofuran, ethyl acetate, acetone, acetonitrile, dimethylsulfoxide, dimethyl-

formamide, or alcohols such as methanol, ethanol, n-propanol, isopropanol, n-butanol or t-butanol, or hydrocarbons such as pentane, hexane, cyclohexane, benzene, toluene or xylene, or halogeno-hydrocarbons such as dichloromethane, dichloroethane, trichloromethane or chlorobenzene. It is also possible to use mixtures of the above-mentioned solvents. Preferred for the process is tetrahydrofuran.

Suitable acids for the process  $(II) + (III)/(VI) + (IV) \rightarrow (IB)$  are generally inorganic or organic acids. These preferably include carboxylic acids, such as, for example, acetic acid or trifluoroacetic acid, sulfonic acids, such as, for example, methanesulfonic acid or p-toluenesulfonic acid, hydrochloric acid or phosphoric acids such as polyphosphoric acids. Preference is given to polyphosphoric acid ethyl ester. The acid is employed in an amount from 0.25 mol to 100 mol, relative to 1 mol of the compound of the general formula (III).

The process is in general carried out in a temperature range from +20°C to +150°C, preferably from +60°C to +100°C.

The process is generally carried out at normal pressure. However, it is also possible to carry it out at elevated pressure or at reduced pressure (for example in a range from 0.5 to 5 bar).

Suitable solvents for the process  $(IB) + (V) \rightarrow (I)$  are generally customary organic solvents which do not change under the reaction conditions. These include ethers such as diethyl ether, diisopropyl ether, 1,2-dimethoxyethane, dioxan or tetrahydrofuran, ethyl acetate, acetone, acetonitrile, dimethylsulfoxide, dimethylformamide, or hydrocarbons such as pentane, hexane, cyclohexane, benzene, toluene or xylene, or halogeno-hydrocarbons such as dichloromethane, dichloroethane, trichloromethane or chlorobenzene. It is also possible to use mixtures of the above-mentioned solvents. Preferred for the process is tetrahydrofuran.



- 20 -

Suitable bases for the process (IB) + (V)  $\rightarrow$  (I) are generally inorganic or organic bases. These preferably include cyclic amines, such as, for example, piperidine or 4-N,N-dimethylaminopyridine, or (C<sub>1</sub>-C<sub>4</sub>)-trialkylamines, such as, for example, triethylamine or diisopropylethylamine, or hydrides such as sodium hydride.

5 Preference is given to sodium hydride. The base is employed in an amount from 0.1 mol to 10 mol, preferably from 1 mol to 3 mol, relative to 1 mol of the compound of general formula (IV).

10 The process is in general carried out in a temperature range from 0°C to +150°C, preferably from +20°C to +80°C, especially at room temperature.


The process is generally carried out at normal pressure. However, it is also possible to carry it out at elevated pressure or at reduced pressure (for example in a range from 0.5 to 5 bar).

15

The compounds of the general formulas (II), (III), (IV), (V) and (VI) are known per se, or they can be prepared by customary methods.

The above-mentioned method can be illustrated by the following scheme:

20



The compounds according to the invention exhibit an unforeseeable, useful pharmacological and pharmacokinetic activity spectrum. They are therefore suitable for use as medicaments for the treatment and/or prophylaxis of disorders in humans and animals.

5

Surprisingly, the compounds of the present invention show human neutrophil elastase (HNE) inhibitory activity and are therefore suitable for the preparation of medicaments for the treatment of diseases associated with HNE activity. They may thus provide an effective treatment of acute and chronic inflammatory processes, such as rheumatoid arthritis, atherosclerosis, and especially of acute and chronic pulmonary diseases, such as lung fibrosis, cystic fibrosis, pneumonia, acute respiratory distress syndrome (ARDS), in particular pulmonary emphysema, including smoking-induced emphysema, and chronic obstructive pulmonary diseases (COPD), chronic bronchitis and bronchiectasis. The compounds of the present invention may further provide an effective treatment for cardiovascular ischaemic diseases such as acute coronary syndrome, acute myocardial infarction, unstable and stable angina pectoris, coronary artery bypass grafts (CABG) and heart failure development, for atherosclerosis, mitral valvular disease, atrial septal defects, percutaneous transluminal coronary angioplasty (PTCA), inflammation after open heart surgery and for pulmonary hypertension. They may also prove useful for an effective treatment of rheumatoid arthritis, acute inflammatory arthritis, cancer, acute pancreatitis, ulcerative colitis, periodontal disease, Chury-Strauss syndrome, acute and chronic atopic dermatitis, psoriasis, systemic lupus erythematosus, bullous pemphigus, sepsis, alcoholic hepatitis, liver fibrosis, Behcet's disease, allergic fungal sinusitis, allergic sinusitis, Crohn's disease, Kawasaki disease, glomerulonephritis, acute pyelonephritis, colorectal diseases, chronic suppurative otitis media, chronic venous leg ulcers, inflammatory bowel disease, bacterial and viral infections, brain trauma, stroke and other conditions in which neutrophil participation is involved.

30

The present invention further provides medicaments containing at least one compound according to the invention, preferably together with one or more pharma-

cologically safe excipient or carrier substances, and also their use for the above-mentioned purposes.

5 The active component can act systemically and/or locally. For this purpose, it can be applied in a suitable manner, for example orally, parenterally, pulmonally, nasally, sublingually, lingually, buccally, rectally, transdermally, conjunctivally, otically or as an implant.

10 For these application routes, the active component can be administered in suitable application forms.

Useful oral application forms include application forms which release the active component rapidly and/or in modified form, such as for example tablets (non-coated and coated tablets, for example with an enteric coating), capsules, sugar-coated  
15 tablets, granules, pellets, powders, emulsions, suspensions, solutions and aerosols.

Parenteral application can be carried out with avoidance of an absorption step (intravenously, intraarterially, intracardially, intraspinally or intralumbally) or with inclusion of an absorption (intramuscularly, subcutaneously, intracutaneously, percutaneously or intraperitoneally). Useful parenteral application forms include  
20 injection and infusion preparations in the form of solutions, suspensions, emulsions, lyophilisates and sterile powders.

Forms suitable for other application routes include for example inhalatory pharmaceutical forms (including powder inhalers, nebulizers), nasal drops/solutions, sprays; tablets or capsules to be administered lingually, sublingually or buccally, suppositories, ear and eye preparations, vaginal capsules, aqueous suspensions (lotions, shake mixtures), lipophilic suspensions, ointments, creams, milk, pastes, dusting powders or implants.

The active components can be converted into the recited application forms in a manner known per se. This is carried out using inert non-toxic, pharmaceutically suitable excipients. These include inter alia carriers (for example microcrystalline cellulose), solvents (for example liquid polyethylene glycols), emulsifiers (for example sodium dodecyl sulphate), dispersing agents (for example polyvinylpyrrolidone), synthetic and natural biopolymers (for example albumin), stabilizers (for example antioxidants such as ascorbic acid), colorants (for example inorganic pigments such as iron oxides) or taste and/or odor corrigents.

For human use, in the case of oral administration, it is recommendable to administer doses of from 0.001 to 50 mg/kg, preferably of 0.01 mg/kg to 20 mg/kg. In the case of parenteral administration, such as, for example, intravenously or via mucous membranes nasally, buccally or inhalationally, it is recommendable to use doses of 0.001 mg/kg to 0.5 mg/kg.

In spite of this, it can be necessary in certain circumstances to depart from the amounts mentioned, namely as a function of body weight, application route, individual behaviour towards the active component, manner of preparation and time or interval at which application takes place. It can for instance be sufficient in some cases to use less than the aforementioned minimum amount, while in other cases the upper limit mentioned will have to be exceeded. In the case of the application of larger amounts, it can be advisable to divide them into a plurality of individual doses spread through the day.

The percentages in the tests and examples which follows are, unless otherwise stated, by weight; parts are by weight. Solvent ratios, dilution ratios and concentrations reported for liquid/liquid solutions are each based on the volume.

### **A. Evaluation of physiological activity**

The potential of the compounds of the invention to inhibit neutrophil elastase activity may be demonstrated, for example, using the following assays:

5

#### **I. *In vitro* enzyme assays of human neutrophil elastase (HNE)**

##### **Assay contents**

10 assay buffer: 0.1 M HEPES-NaOH buffer pH 7.4, 0.5 M NaCl, 0.1% (w/v) bovine serum albumin;

suitable concentration (see below) of HNE (18 U/mg lyophil., #20927.01, SERVA Electrophoresis GmbH, Heidelberg, Germany) in assay buffer;

15

suitable concentration (see below) of substrate in assay buffer;

suitable concentration of test compounds diluted with assay buffer from a 10 mM stock solution in DMSO.

20

##### **Example A**

***In vitro* inhibition of HNE using a fluorogenic peptide substrate (continuous read-out signal, 384 MTP assay format):**

25

In this protocol, the elastase substrate MeOSuc-Ala-Ala-Pro-Val-AMC (#324740, Calbiochem-Novabiochem Corporation, Merck KGaA, Darmstadt, Germany) is used. The test solution is prepared by mixing 10  $\mu$ l of test compound dilution, 20  $\mu$ l of HNE enzyme dilution (final concentration 8 - 0.4  $\mu$ U/ml, routinely 2.1  $\mu$ U/ml) and  
30 20  $\mu$ l of substrate dilution (final concentration 1 mM - 1  $\mu$ M, routinely 20  $\mu$ M), respectively. The solution is incubated for 0 - 2 hrs at 37°C (routinely one hour). The

fluorescence of the liberated AMC due to the enzymatic reaction is measured at 37°C (TECAN spectra fluor plus plate reader). The rate of increase of the fluorescence (ex. 395 nm, em. 460 nm) is proportional to elastase activity. IC<sub>50</sub> values are determined by RFU-versus-[I] plots. K<sub>m</sub> and K<sub>m(app.)</sub> values are determined by Lineweaver-Burk  
 5 plots and converted to K<sub>i</sub> values by Dixon plots.

The preparation examples had IC<sub>50</sub> values within the range of 5 nM - 5 µM in this assay. Representative data are given in Table 1:

10 Table 1

Example No.	IC <sub>50</sub> [nM]
1	8
9	40
14	5
15	8
16	10
20	700
24	13
26	10
28	50
58	1100
60	5
72	6
73	60
74	20
103	60
109	15
110	50

### Example B

***In vitro* inhibition of HNE using a fluorogenic, insoluble elastin substrate (discontinuous read-out signal, 96 MTP assay format):**

5  
In this protocol the elastase substrate elastin-fluorescein (#100620, ICN Biomedicals GmbH, Eschwege, Germany) is used. The test solution is prepared by mixing 3 µl of test compound dilution, 77 µl of HNE enzyme dilution (final concentration 0.22 U/ml - 2.2 mU/ml, routinely 21.7 µU/ml) and 80 µl substrate suspension (final  
10 concentration 2 mg/ml). The suspension is incubated for 0 - 16 hrs at 37°C (routinely four hours) under slightly shaking conditions. To stop the enzymatic reaction, 160 µl of 0.1 M acetic acid are added to the test solution (final concentration 50 mM). The polymeric elastin-fluorescein is pulled down by centrifugation (Eppendorf 5804 centrifuge, 3.000 rpm, 10 min). The supernatant is transferred into a new MTP and  
15 the fluorescence of the liberated peptide fluorescein due to the enzymatic reaction is measured (BMG Fluostar plate reader). The rate of fluorescence (ex. 490 nm, em. 520 nm) is proportional to elastase activity. IC<sub>50</sub> values are determined by RFU-versus-[I] plots.

## 20 II. *In vitro* human neutrophil assays

### Example A

***In vitro* PMN elastolysis assay:**

25  
This assay is used to determine the elastolytic potential of human polymorphonuclear cells (PMNs) and assess the proportion of degradation due to neutrophil elastase [cf. Z.W. She et al., Am. J. Respir. Cell. Mol. Biol. 9, 386-392 (1993)].

30  
Tritiated elastin, in suspension, is coated on to a 96 well plate at 10 µg per well. Test and reference [ZD-0892 (J. Med. Chem. 40, 1876-1885, 3173-3181 (1997), WO

95/21855) and  $\alpha$ 1 protease inhibitor ( $\alpha$ 1PI)] compounds are added to the wells at the appropriate concentrations. Human PMNs are separated from peripheral venous blood of healthy donors and resuspended in culture media. The neutrophils are added to the coated wells at concentrations ranging between  $1 \times 10^6$  to  $1 \times 10^5$  cells per well. Porcine pancreatic elastase ( $1.3 \mu\text{M}$ ) is used as a positive control for the assay, and  $\alpha$ 1PI ( $1.2 \mu\text{M}$ ) is used as the positive inhibitor of neutrophil elastase. The cellular control is PMNs without compound at each appropriate cell density. The cells plus compounds are incubated in a humidified incubator at  $37^\circ\text{C}$  for 4 hours. The plates are centrifuged to allow the harvest of cell supernatant only. The supernatant is transferred in  $75 \mu\text{l}$  volumes to corresponding wells of a 96 well Lumaplate™ (solid scintillant containing plates). The plates are dried until no liquid is visible in the wells and read in a beta counter for 3 minutes per well.

Elastolysis of the  $^3\text{H}$ -elastin results in an increase in counts in the supernatant. An inhibition of this elastolysis shows a decrease, from the cellular control, of tritium in the supernatant.  $\alpha$ 1PI gave  $83.46 \pm 3.97\%$  (mean  $\pm$  s.e.m.) inhibition at  $1.2 \mu\text{M}$  ( $n = 3$  different donors at  $3.6 \times 10^5$  cells per well).  $\text{IC}_{50}$  values were obtained for the reference compound ZD-0892 of  $45.50 \pm 7.75 \text{ nM}$  (mean  $\pm$  s.e.m.) ( $n = 2$  different donors at  $3.6 \times 10^5$  cells per well).

Given that ZD-0892 is a selective inhibitor of PMN elastase along with the data from  $\alpha$ 1PI inhibition, these results indicate that the majority of elastin degradation by PMNs is due to the release of neutrophil elastase, and not to another elastolytic enzyme such as matrix metalloproteases (MMPs). The compounds of this invention are evaluated for their inhibitory activity in this HNE-dependent model of neutrophil elastolysis.



### **Example B**

#### ***In vitro* inhibition of membrane bound elastase:**

5 Measurement of the inhibition of elastase bound to neutrophil membranes is performed using a human neutrophil assay. Neutrophils are stimulated with LPS at 37°C for 35 min and then spun at 1600 rpm. Subsequently, the membrane bound elastase is fixed to the neutrophils with 3% paraformaldehyde and 0.25% glutaraldehyde for 3 min at 4°C. The neutrophils are then spun, and vehicle and the compound under  
10 evaluation are added, followed by addition of the substrate MeOSuc-Ala-Ala-Pro-Val-AMC (#324740, Calbiochem-Novabiochem Corporation, Merck KGaA, Darmstadt, Germany) at 200 µM. Following a 25 min incubation at 37°C, the reaction is terminated with PMSF (phenylmethanesulfonyl fluoride), and the fluorescence is read at ex: 400 nm and em: 505 nm. IC<sub>50</sub> values are determined by inter-  
15 polation from plots of relative fluorescence vs. inhibitor concentration.

### **III. *In vivo* models**

#### **Example A**

20

#### ***In vivo* model of acute lung injury in the rat:**

Instillation of human neutrophil elastase (HNE) into rat lung causes acute lung damage. The extent of this injury can be assessed by measuring lung haemorrhage.

25

Rats are anaesthetised with Hypnorm/Hypnovel/water and instilled with HNE or saline delivered by microsyringe into the lungs. Test compounds are administered by intravenous injection, by oral gavage or by inhalation at set times prior to the administration of HNE. Sixty minutes after the administration of elastase animals are  
30 killed by an anaesthetic overdose (sodium pentobarbitone) and the lungs lavaged with 2 ml heparinised phosphate buffered saline (PBS). Bronchoalveolar lavage

(BAL) volume is recorded and the samples kept on ice. Each BAL sample is centrifuged at 900 r.p.m. for 10 minutes at 4-10°C. The supernatant is discarded and the cell pellet resuspended in PBS and the sample spun down again. The supernatant is again discarded and the cell pellet resuspended in 1 ml 0.1% cetyltrimethylammonium bromide (CTAB) / PBS to lyse the cells. Samples are frozen until blood content is assayed. Prior to the haemorrhage assay the samples are defrosted and mixed. 100 µl of each sample are placed into a separate well of a 96 well flat-bottomed plate. All samples are tested in duplicate. 100 µl 0.1% CTAB/PBS is included as a blank. The absorbance of the well contents is measured at 415 nm using a spectrophotometer. A standard curve is constructed by measuring the OD at 415 nm of different concentrations of blood in 0.1% CTAB/PBS. Blood content values are calculated by comparison to the standard curve (included in each plate) and normalised for the volume of BAL fluid retrieved.

The compounds of this invention are evaluated intravenously, orally or by inhalation for their inhibitory activity in this model of HNE-induced haemorrhage in the rat.

### **Example B**

#### ***In vivo* model of acute myocardial infarction in the rat:**

Elastase inhibitors are tested in a rat thread infarct model. Male Wistar rats (weighing >300 g) receive 10 mg/kg aspirin 30 min prior to surgery. They are anaesthetized by isofluran and ventilated (120-130 strokes/min, 200-250 µl stroke volume; MiniVent Type 845, Hugo Sachs Elektronik, Germany) during the whole surgery. Following a left thoracotomy at the fourth intercostal space, the pericardium is opened and the heart briefly exteriorized. A thread is turned around the left coronary artery (LAD) without occluding the artery. The thread is passed under the skin to the neck of the animal. The thorax is closed and the animal is allowed to recover for 4 days. At the fifth day, rats are anaesthetized with ether for 3 min, and the thread is tied and the LAD occluded under ECG control. Test compounds are administered before or after

LAD occlusion per os, intraperitoneally or intravenously (bolus or permanent infusion). After 1 hr occlusion, the thread is reopened to allow reperfusion. Hearts are excised, and infarct sizes are determined 48 hours later by staining of the re-occluded hearts with Evans blue, followed by TTC (triphenyltetrazolium chloride) staining of  
5 2 mm heart sections. Normoxic (not occluded tissue) areas stain blue, ischemic (occluded but surviving tissue) areas stain red and necrotic (occluded dead tissue) areas remain white. Each tissue section is scanned and infarct sizes are determined by computer planimetry.

10 **B. Examples**

**Abbreviations:**

aq.	aqueous
conc.	concentrated
DMF	N,N-dimethylformamide
DMSO	dimethylsulfoxide
EI	electron impact ionisation (for MS)
ESI	electro-spray ionisation (for MS)
HPLC	high pressure liquid chromatography
LC-MS	liquid chromatography coupled with mass spectroscopy
Mp.	melting point
MS	mass spectroscopy
NMR	nuclear magnetic resonance spectroscopy
of th.	of theoretical (yield)
R <sub>t</sub>	retention time (for HPLC)
THF	tetrahydrofuran

**General methods:**

All reactions are carried out under an argon atmosphere unless otherwise noted. Solvents are used as purchased from Aldrich without further purification. 'Silica gel' or 'Silica' refers to Silica gel 60 (0.040 mm-0.063 mm) from Merck KGaA company. Melting points were obtained with a Büchi 512 or similar melting point device and are uncorrected.

Compounds purified by preparative HPLC are purified over a RP18-column with acetonitrile and water as the eluent, using a 1:9 to 9:1 gradient.

**LC-MS / HPLC methods:****LC-MS method 1**

Instrument: Micromass Quattro LCZ, HP1100; column: Uptisphere HDO, 50 mm x 2.0 mm, 3 µm; eluent A: water + 0.05% formic acid, eluent B: acetonitrile + 0.05% formic acid; gradient: 0.0 min 100% A → 0.2 min 100% A → 2.9 min 30% A → 3.1 min 10% A → 4.5 min 10% A; oven: 55°C; flow: 0.8 ml/min; UV-detection: 208-400 nm.

**LC-MS method 2**

Instrument: Waters Alliance 2790 LC; column: Symmetry C18, 50 mm x 2.1 mm, 3.5 µm; eluent A: water + 0.1% formic acid, eluent B: acetonitrile + 0.1% formic acid; gradient: 0.0 min 5% B → 5.0 min 10% B → 6.0 min 10% B; temperature: 50°C; flow: 1.0 ml/min; UV-detection: 210 nm.

**LC-MS method 3**

Instrument: Micromass Platform LCZ, HP1100; column: Aquasil C-18, 50 mm x 2.0 mm, 3 µm; eluent A: water + 0.05% formic acid, eluent B: acetonitrile + 0.05% formic acid; gradient: 0.0 min 100% A → 0.2 min 100% A → 2.9 min 30% A → 3.1

min 10% A → 4.5 min 10% A; oven: 55°C; flow: 0.8 ml/min; UV-detection: 208-400 nm.

#### HPLC method 4

- 5 Instrument: HP 1100 with DAD-detection; column: Kromasil RP-18, 60 mm x 2 mm, 3.5 µm; eluent: A = 5 ml HClO<sub>4</sub>/l H<sub>2</sub>O, B = acetonitrile; gradient: 0 min 2% B, 0.5 min 2% B, 4.5 min 90% B, 6.5 min 90% B; flow: 0.75 ml/min; temperature: 30°C; UV-detection: 210 nm.

#### 10 LC-MS method 5

- Instrument: Micromass TOF-MUX-Interface 4-fold parallel injection, with HPLC Waters 600; column: Uptisphere HDO, 50 mm x 2.0 mm, 3.0 µm; eluent A: 1 l water + 1 ml 50% formic acid, eluent B: 1 l acetonitrile + 1 ml 50% formic acid; gradient: 0.0 min 100% A → 0.2 min 100% A → 2.9 min 30% A → 3.1 min 10% A → 4.5 min 10% A → 4.6 min 100% A → 6.5 min 100% A; oven: room temperature; flow: 0.8 ml/min; UV-detection: 210 nm.

#### LC-MS method 6


- Instrument: Micromass Platform LCZ with HPLC Agilent Serie 1100; column: Grom-SIL120 ODS-4 HE, 50 mm x 2.0 mm, 3 µm; eluent A: 1 l water + 1 ml 50% formic acid, eluent B: 1 l acetonitrile + 1 ml 50% formic acid; gradient: 0.0 min 100% A → 0.2 min 100% A → 2.9 min 30% A → 3.1 min 10% A → 4.5 min 10% A; oven: 55°C; flow: 0.8 ml/min; UV-detection: 208-400 nm.

#### 25 LC-MS method 7

- Instrument: Micromass Quattro LCZ with HPLC Agilent Serie 1100; column: Uptisphere HDO, 50 mm x 2.0 mm, 3 µm; eluent A: 1 l water + 1 ml 50% formic acid, eluent B: 1 l acetonitrile + 1 ml 50% formic acid; gradient: 0.0 min 100% A → 0.2 min 100% A → 2.9 min 30% A → 3.1 min 10% A → 4.5 min 10% A; oven: 55°C; flow: 0.8 ml/min; UV-detection: 208-400 nm.

**Starting Materials:****Example 1A****2-Bromo-5-(1,3-dioxolan-2-yl)pyridine**

5



6-Bromo-3-pyridinecarbaldehyde (500 mg, 2.7 mmol) and 1,2-ethanediol (200 mg, 3.2 mmol) are dissolved in toluene (50 ml) together with Amberlyst 15 (100 mg) in a round bottom flask equipped with a reflux condenser and a Dean-Stark trap. The solution is stirred at reflux overnight, then cooled to room temperature, filtered and concentrated *in vacuo*. The crude product is chromatographed over silica gel with cyclohexane and ethyl acetate as the eluent to afford the title compound as a colorless oil.


15 Yield: 0.489 g (79% of th.)

HPLC (method 4): 3.46 min.

MS (ESIpos):  $m/z = 231$  ( $M+H$ )<sup>+</sup>

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  = 8.46 (d, 1H), 7.64 (m, 1H), 7.49 (m, 1H), 4.15-4.00 (m, 4H) ppm.

20

**Example 2A****5-(1,3-Dioxolan-2-yl)-2-pyridinecarbonitrile**

5


Example 1A (2.8 g, 12.5 mmol), zinc cyanide (1.6 g, 13.8 mmol) and tetrakis-(triphenylphosphine)palladium(0) (1.4 g, 1.3 mmol) are dissolved in dimethylformamide (100 ml) and stirred overnight (18 h) at 80°C. Additional tetrakis-(triphenylphosphine)palladium(0) (0.1 g) is added and the reaction is stirred again  
10 overnight (18 h) at 80°C, then allowed to stand at room temperature for 2 days (48 hours). The solvent is removed *in vacuo*, to the residue is given water (100 ml) and the product is extracted with ethyl acetate (1 l). The organic phase is washed with brine (200 ml), dried with magnesium sulphate monohydrate, filtered and concentrated *in vacuo*. The crude product is chromatographed over silica gel with  
15 cyclohexane and ethyl acetate as the eluent to afford the title compound as a white amorphous solid.

Yield: 0.94 g (42% of th.)

HPLC (method 4): 3.21 min.

MS (ESIpos):  $m/z = 177 (M+H)^+$

20  $^1\text{H-NMR}$  (400 MHz, DMSO- $d_6$ ):  $\delta = 8.81$  (s, 1H), 8.09 (s, 2H), 5.95 (s, 1H), 4.13-3.94 (m, 4 H) ppm.

**Example 3A****5-Formyl-2-pyridinecarbonitrile**

5

*Method a):*

Prepared in analogy to the procedure of Dodd, D. *et al.* [*J.Org.Chem.* **1992**, *57*, 7226-7234]: To a stirred solution of 5-(1,3-dioxolan-2-yl)-2-pyridinecarbonitrile (Example 2A; 850 mg, 4.8 mmol) in acetone/water 85:15 (59.5 ml) is given *p*-toluenesulphonic acid (102 mg, 0.59 mmol). The reaction is stirred at reflux overnight (18 h), then additional *p*-toluenesulphonic acid (50 mg) and water (5 ml) are added. The reaction is stirred at reflux for an additional 48 h. The solution is cooled to room temperature and quenched with saturated sodium bicarbonate solution. The product is extracted with ethyl acetate (3 x 100 ml), dried over magnesium sulphate monohydrate, filtered and concentrated *in vacuo*. The crude product is purified by preparative HPLC to afford a pale yellow solid.

15

Yield: 0.66 g (93% of th.)

Mp.: 80-82°C

HPLC (method 4): 2.13 min.

20 MS (ESIpos):  $m/z = 133$  (M+H)<sup>+</sup>

<sup>1</sup>H-NMR (400 MHz, DMSO-d<sub>6</sub>):  $\delta = 10.18$  (s, 1H), 9.21 (m, 1H), 8.49 (m, 1H), 8.27 (m, 1H) ppm.

*Method b):*

25 1.04 g (8.2 mmol) oxalylchloride are dissolved in 8 ml dichloromethane. At -78°C, 1.28 g (16.4 mmol) dimethylsulfoxide are added dropwise. The solution is stirred at -78°C for 20 minutes, then 1 g (7.46 mmol) of the compound of Example 5A,



- 36 -

dissolved in 7 ml dichloromethane, is added, and stirring at  $-78^{\circ}\text{C}$  is continued for another 2 hours. 3.4 g (33.6 mmol) triethylamine are then added dropwise, and after warming up to room temperature, the mixture is purified by column chromatography (silica, eluent cyclohexane to cyclohexane/ethyl acetate 2:1).


5 Yield: 0.76 g (77% of th.)

Analytical data: see above.

#### Example 4A

5-Methyl-2-pyridinecarbonitrile


10



36 g (209 mmol) 2-bromo-5-methylpyridine and 37.5 g (418 mmol) copper cyanide are refluxed for two hours in 500 ml dimethylformamide. After cooling down to  $50^{\circ}\text{C}$ , 10% aqueous ammonia solution (500 ml) is added with stirring. The product is extracted with dichloromethane, the organic phase is dried over magnesium sulfate, and the solvent is removed *in vacuo*. The product is purified by column chromatography (silica, eluent cyclohexane/ethyl acetate 9:1):

15 Yield: 18 g (73% of th.)

20  $^1\text{H-NMR}$  (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  = 2.4 (s, 3H), 7.6 (m, 2H), 8.6 (s, 1H) ppm.

**Example 5A****5-(Hydroxymethyl)-2-pyridinecarbonitrile**

5

The compound of Example 4A (13 g, 110 mmol) is dissolved in 400 ml tetrachloromethane, and 29.4 g (165 mmol) N-bromosuccinimide and 0.4 g (1.6 mmol) dibenzoylperoxide are added. The reaction mixture is refluxed for three hours, cooled down to room temperature and filtered. The solution is washed with aqueous sodium thiosulfate, dried over magnesium sulfate, and the solvent is removed *in vacuo*. The residue is dissolved in 200 ml dioxane and 200 ml water, calciumcarbonate (44 g, 440 mmol) is added, and the mixture is stirred at reflux for 2 hours. After cooling down to room temperature, the mixture is filtered, and dichloromethane is added. After phase separation, the organic phase is dried over magnesium sulfate, and the solvent is removed *in vacuo*. The product is purified by chromatography (silica, eluent cyclohexane/ethyl acetate 2:1).

10

15


Yield: 5.2 g (35% of th.)

<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ = 4.7 (d, 2H), 5.6 (t, 1H), 8.0 (m, 2H), 8.7 (s, 1H) ppm.

20

**Preparation Examples:****Example 1**

5 Ethyl 4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate




10 7.0 g (34.29 mmol) N-[3-(trifluoromethyl)phenyl]urea, 8.99 g (68.58 mmol) 4-cyanobenzaldehyde, 8.92 g (68.58 mmol) ethyl 3-oxobutanoate and 20 g polyphosphoric acid ethyl ester are suspended in 250 ml of THF. The mixture is stirred at reflux for 18 hours. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with cyclohexane/ethyl acetate as eluent.

15 Yield: 13.4 g (91%)

<sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>): δ = 1.1 (t, 3H); 2.0 (s, 3H); 4.0 (q, 2H); 5.4 (d, 1H); 7.6 (m, 3H); 7.7 (m, 3H); 7.9 (m, 2H); 8.4 (d, 1H) ppm.

**Example 2**

4-{5-Acetyl-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-4-pyrimidinyl}benzonitrile



5

265 mg (1.3 mmol) N-[3-(trifluoromethyl)phenyl]urea, 131 mg (1.0 mmol) 4-cyano-benzaldehyde, and 100 mg (1.0 mmol) 2,4-pentanedione are suspended in 2 ml of THF, and catalytic amounts of concentrated hydrochloric acid are added. The mixture is stirred at reflux for 18 hours. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with cyclohexane/ethyl acetate as eluent.

10


Yield: 29 mg (7%)

<sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>): δ = 2.0 (s, 3H); 2.2 (s, 3H); 5.5 (d, 1H); 7.5 (m, 1H); 7.6 (m, 3H); 7.7 (m, 1H); 7.8 (m, 1H); 7.9 (m, 2H); 8.5 (d, 1H) ppm.

15

**Example 3**

Ethyl 4-(4-bromophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

204 mg (1.0 mmol) N-[3-(trifluoromethyl)phenyl]urea, 142 mg (0.77 mmol) 4-bromobenzaldehyde, and 100 mg (0.77 mmol) ethyl 3-oxobutanoate are suspended in 2 ml of THF, and catalytic amounts of concentrated hydrochloric acid are added. The mixture is stirred at reflux for 18 hours. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with cyclohexane/ethyl acetate as eluent.

10


Yield: 23 mg (6%)

15

$^1\text{H-NMR}$  (200 MHz, DMSO- $d_6$ ):  $\delta$  = 1.1 (t, 3H); 2.0 (s, 3H); 4.0 (q, 2H); 5.3 (d, 1H); 7.4 (m, 2H); 7.6 (m, 3H); 7.7 (m, 3H); 8.3 (d, 1H) ppm.

**Example 4**

Ethyl 4-(4-cyanophenyl)-6-methyl-2-oxo-1-[4-fluorophenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

154 mg (1.0 mmol) N-[4-fluorophenyl]urea, 101 mg (0.77 mmol) 4-cyanobenzaldehyde, and 100 mg (0.77 mmol) ethyl 3-oxobutanoate are suspended in 2 ml of THF, and catalytic amounts of concentrated hydrochloric acid are added. The mixture is stirred at reflux for 18 hours. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with cyclohexane/ethyl acetate as eluent.

10


Yield: 40 mg (14%)

<sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>): δ = 1.1 (t, 3H); 2.0 (s, 3H); 4.0 (q, 2H); 5.3 (d, 1H); 7.3 (m, 4H); 7.5 (m, 2H); 7.9 (m, 2H); 8.3 (d, 1H) ppm.

15

**Example 5**

Ethyl 4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-chlorophenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

170 mg (1.0 mmol) N-[3-chlorophenyl]urea, 100 mg (0.77 mmol) 4-cyanobenzaldehyde and 100 mg (0.77 mmol) ethyl 3-oxobutanoate are suspended in 2 ml of THF, and catalytic amounts of concentrated hydrochloric acid are added. The mixture is stirred at reflux for 18 hours. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with cyclohexane/ethyl acetate as eluent.


Yield: 13 mg (4%)

<sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>): δ = 1.1 (t, 3H); 2.1 (s, 3H); 4.0 (q, 2H); 5.3 (d, 1H); 7.2 (m, 1H); 7.4 (m, 3H); 7.5 (m, 2H); 7.9 (m, 2H); 8.3 (d, 1H) ppm.

15

**Example 6**

(1S)-2-Methoxy-1-methyl-2-oxoethyl 4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

200 mg (0.98 mmol) N-[3-(trifluoromethyl)phenyl]urea, 129 mg (0.98 mmol) 4-cyanobenzaldehyde, 92 mg (0.49 mmol) (1S)-2-methoxy-1-methyl-2-oxoethyl 3-oxobutanoate, and 295 mg polyphosphoric acid ethyl ester are suspended in 3 ml of THF. The mixture is stirred at reflux for 18 hours. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with cyclohexane/ethyl acetate as eluent. A mixture of diastereoisomers is obtained.

10

Yield: 96 mg (40%)


15

<sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>): δ = 1.3 (d, 3H); 1.4 (d, 3H); 2.0 (s, 3H+3H); 3.6 (s, 3H); 3.6 (s, 3H); 5.0 (m, 1H+1H); 5.4 (m, 1H+1H); 7.6-7.9 (m, 8H+8H); 8.4 (m, 1H+1H) ppm.



**Example 7**

4-{6-Methyl-5-(4-morpholinylcarbonyl)-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-4-pyrimidinyl} benzonitrile



5

150 mg (0.73 mmol) N-[3-(trifluoromethyl)phenyl]urea, 96 mg (0.73 mmol) 4-cyanobenzaldehyde, 63 mg (0.37 mmol) 4-(4-morpholinyl)-4-oxo-2-butanone and 220 mg polyphosphoric acid ethyl ester are suspended in 3 ml of THF. The mixture is stirred at reflux for 18 hours. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with dichloromethane/methanol as eluent.

10


Yield: 28 mg (16%)

<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ = 1.5 (s, 3H); 3.1 (m, 4H); 3.6 (m, 4H); 5.3 (br.s, 1H); 7.6 (m, 2H); 7.7 (m, 1H); 7.8 (m, 2H); 7.9 (m, 2H); 8.0 (br.s, 1H) ppm.

15

**Example 8**

4-(4-Cyanophenyl)-N,N-diethyl-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-  
1,2,3,4-tetrahydro-5-pyrimidinecarboxamide



5

200 mg (0.98 mmol) N-[3-(trifluoromethyl)phenyl]urea, 128 mg (0.98 mmol) 4-cyanobenzaldehyde, 77 mg (0.49 mmol) 4-(4-diethylamino)-4-oxo-2-butanone and 295 mg polyphosphoric acid ethyl ester are suspended in 3 ml of THF. The mixture is stirred at reflux for 18 hours. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with dichloromethane/methanol as eluent.

10


Yield: 106 mg (47%)

15

<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ = 0.9 (m, 6H); 3.1 (m, 4H); 5.2 (br.s, 1H); 7.6 (m, 2H); 7.7 (m, 1H); 7.8 (m, 2H); 7.9 (m, 2H); 8.0 (br.s, 1H) ppm.

**Example 9**

6-Amino-4-(4-cyanophenyl)-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarbonitrile



5

400 mg (1.97 mmol) N-[3-(trifluoromethyl)phenyl]urea, 199 mg (1.51 mmol) 4-cyanobenzaldehyde and 100 mg (1.51 mmol) malononitrile are suspended in 2 ml of THF, and catalytic amounts of concentrated hydrochloric acid are added. The mixture is stirred at reflux for 18 hours. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with dichloromethane/methanol as eluent.

10


Yield: 4 mg (1%)

<sup>1</sup>H-NMR (400 MHz, DMSO-d<sub>6</sub>): δ = 5.2 (d, 1H); 6.0 (s, 2H); 7.6 (m, 3H); 7.7 (m, 2H); 7.8 (m, 1H); 7.9 (m, 2H) 8.4 (d, 1H) ppm.

15

**Example 10**

Ethyl 4-(4-cyanophenyl)-3-formyl-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

100 mg (0.23 mmol) of Example 1 are dissolved in 1 ml dimethylformamide, and 35.7 mg (0.23 mmol) phosphorylchloride are added. The reaction mixture is stirred at 70°C for two hours. After cooling down to room temperature, the product is isolated by preparative HPLC.


10

Yield: 43 mg (41%)

<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ = 1.1 (t, 3H); 2.1 (s, 3H); 4.1 (q, 2H); 6.4 (s, 1H); 7.6 (m, 2H); 7.7 (m, 1H); 7.8 (m, 1H); 7.9 (m, 4H); 9.2 (s, 1H) ppm.

**Example 11**

4-(4-Cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylic acid



5

3 g (7 mmol) of Example 1 are dissolved in a mixture of 50 ml water and 100 ml 5% KOH in ethanol. The reaction mixture is stirred at room temperature for 18 hours. The solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with dichloromethane/methanol as eluent.


10

Yield: 1.27 g (45%)

<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ = 2.0 (s, 3H); 5.4 (d, 1H); 7.6 (m, 1H); 7.6 (m, 2H); 7.7 (m, 1H); 7.8 (m, 1H); 7.9 (m, 3H); 8.3 (d, 1H); 12.5 (s, 1H) ppm.

**Example 12**

4-(4-Cyanophenyl)-6-methyl-2-oxo-N-propyl-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxamide



5

40 mg (0.1 mmol) of Example 11 are dissolved in 2 ml dimethylformamide, 7 mg (0.11 mmol) n-propylamine, 15 mg (0.11 mmol) 1-hydroxy-1H-benzotriazole hydrate and 12 mg (0.1 mmol) 4-dimethylaminopyridine are added. The reaction mixture is stirred at 0°C, then 21 mg (0.11 mmol) 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride are added. The reaction mixture is stirred at room temperature for 18 hours, then water and ethyl acetate are added. The organic phase is washed with saturated aqueous KHSO<sub>4</sub>, water and brine, dried over sodium sulfate and evaporated to dryness *in vacuo*. If necessary, the product is further purified by column chromatography or preparative HPLC.

15


Yield: 29 mg (66%)

<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ = 0.7 (t, 3H); 1.3 (sext, 2H); 1.7 (s, 3H); 3.0 (q, 2H); 5.4 (d, 1H); 7.6 (m, 3H); 7.7 (m, 2H); 7.8 (m, 2H); 7.9 (m, 1H); 8.1 (d, 1H) ppm.

20

**Example 13**

4-(4-Cyanophenyl)-N-(2-methoxyethyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)-phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxamide



5

48 mg (0.12 mmol) of Example 11 are dissolved in 2 ml dimethylformamide, 10 mg (0.13 mmol) 2-methoxyethylamine, 18 mg (0.13 mmol) 1-hydroxy-1H-benzotriazole hydrate and 15 mg (0.12 mmol) 4-dimethylaminopyridine are added. The reaction mixture is stirred at 0°C, then 25 mg (0.13 mmol) 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride are added. The reaction mixture is stirred at room temperature for 18 hours, then water and ethyl acetate are added. The organic phase is washed with saturated aqueous KHSO<sub>4</sub>, water and brine, dried over sodium sulfate and evaporated to dryness *in vacuo*. If necessary, the product is further purified by column chromatography or preparative HPLC.


15

Yield: 22 mg (40%)

<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ = 1.7 (s, 3H); 3.2 (s, 3H); 3.3 (m, 4H); 5.4 (d, 1H); 7.6 (m, 3H); 7.7 (m, 3H); 7.9 (m, 2H); 8.1 (m, 1H) ppm.

**Example 14**

Ethyl 4-(4-cyanophenyl)-3,6-dimethyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

89 mg (0.21 mmol) of Example 1 are added to a suspension of 12.4 mg (0.31 mmol) 60% sodium hydride (in mineral oil) in 2 ml THF. The mixture is stirred at room temperature for two hours. Then 26 mg (0.21 mmol) dimethylsulfate are added, and the mixture is stirred at room temperature for another 2 hours. Then water and ethyl acetate are added, and the organic phase is washed with water and brine, dried over sodium sulfate and evaporated *in vacuo*. If necessary, the product is further purified by column chromatography or preparative HPLC.

Yield: 85 mg (93%)


<sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>): δ = 1.1 (t, 3H); 2.0 (s, 3H); 2.8 (s, 3H); 4.0 (q, 2H); 5.5 (s, 1H); 7.6 (m, 3H); 7.7 (m, 1H); 7.8 (m, 2H); 7.9 (m, 2H) ppm.

15



**Example 15**

Ethyl 3-acetyl-4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5


100 mg (0.23 mmol) of Example 1 are added to a suspension of 12 mg (0.28 mmol) 60% sodium hydride (in mineral oil) in 2 ml THF. The mixture is stirred at room temperature for two hours. Then 91 mg (1.16 mmol) acetylchloride are added, and the mixture is stirred at room temperature for another 2 hours. Then water and ethyl acetate are added, and the organic phase is washed with water and brine, dried over sodium sulfate and evaporated to dryness *in vacuo*. If necessary, the product is further purified by column chromatography or preparative HPLC.

Yield: 93 mg (85%)

15 <sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>): δ = 1.2 (t, 3H); 2.1 (s, 3H); 2.5 (s, 3H); 4.2 (m, 2H); 6.7 (s, 1H); 7.4 (m, 1H); 7.5 (m, 2H); 7.6 (m, 1H); 7.7 (m, 1H); 7.8 (m, 1H); 7.9 (m, 2H) ppm.

**Example 16**

Diethyl 6-(4-cyanophenyl)-4-methyl-2-oxo-3-[3-(trifluoromethyl)phenyl]-3,6-dihydro-1,5(2H)-pyrimidinedicarboxylate



5


100 mg (0.23 mmol) of Example 1 are added to a suspension of 12 mg (0.28 mmol) 60% sodium hydride (in mineral oil) in 2 ml THF. The mixture is stirred at room temperature for two hours. Then 126 mg (1.16 mmol) ethyl chloridocarbonate are added, and the mixture is stirred at room temperature for another 2 hours. Then water and ethyl acetate are added, and the organic phase is washed with water and brine, dried over sodium sulfate and evaporated to dryness *in vacuo*. If necessary, the product is further purified by column chromatography or preparative HPLC.

Yield: 92 mg (79%)

15 <sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>): δ = 1.2 (t, 3H; t, 3H); 2.1 (s, 3H); 4.2 (m, 2H); 4.3 (q, 2H); 6.4 (s, 1H); 7.4 (m, 1H); 7.5 (m, 3H); 7.7 (m, 1H); 7.8 (m, 1H); 7.9 (m, 2H) ppm.

**Example 17**

Ethyl 4-(4-cyanophenyl)-6-methyl-1-(3-methylphenyl)-2-oxo-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

150 mg (1.0 mmol) N-[3-methylphenyl]urea, 101 mg (0.77 mmol) 4-cyanobenzaldehyde and 100 mg (0.77 mmol) ethyl 3-oxobutanoate are suspended in 2 ml of THF, and catalytic amounts of concentrated hydrochloric acid are added. The mixture is stirred at reflux for 18 hours. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with cyclohexane/ethyl acetate as eluent.


Yield: 8 mg (3%)

<sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>): δ = 1.1 (t, 3H); 2.0 (s, 3H); 2.3 (s, 3H); 4.0 (q, 2H); 5.3 (d, 1H); 7.0 (m, 2H); 7.2 (m, 1H); 7.3 (m, 1H); 7.6 (m, 2H); 7.9 (m, 2H); 8.2 (d, 1H) ppm.

15

**Example 18**

Ethyl 4-(4-chlorophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

204 mg (1.0 mmol) N-[3-(trifluoromethyl)phenyl]urea, 108 mg (0.77 mmol) 4-chlorobenzaldehyde and 100 mg (0.77 mmol) ethyl 3-oxobutanoate are suspended in 2 ml of THF, and catalytic amounts of concentrated hydrochloric acid are added. The mixture is stirred at reflux for 18 hours. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with cyclohexane/ethyl acetate as eluent.

10


Yield: 29 mg (9%)

15

<sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>): δ = 1.1 (t, 3H); 2.0 (s, 3H); 4.0 (q, 2H); 5.3 (d, 1H); 7.5 (m, 5H); 7.6 (m, 1H); 7.7 (m, 2H); 8.3 (d, 1H) ppm.

**Example 19**

Ethyl 6-(bromomethyl)-4-(4-cyanophenyl)-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

3 g (7 mmol) of Example 1 are dissolved in 100 ml chloroform. At 0°C, 558 mg (3.48 mmol) bromine are added dropwise. The mixture is stirred at room temperature for two hours, then the solvent is removed *in vacuo*. The residue is purified by column chromatography on silica with cyclohexane/ethyl acetate as eluent.


10

Yield: 3.2 g (90%)

<sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>): δ = 1.1 (t, 3H); 4.0 (q, 2H, d, 1H); 4.6 (br d, 1H); 5.4 (d, 1H); 7.6 (m, 3H); 7.7 (m, 2H); 7.8 (m, 1H); 7.9 (m, 2H); 8.6 (d, 1H) ppm.

**Example 20**

Ethyl 4-(4-cyanophenyl)-6-[(diethylamino)methyl]-2-oxo-1-[3-(trifluoromethyl)-phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

20 mg (0.04 mmol) of Example 19 are dissolved in 2 ml acetone, and 8 mg (0.10 mmol) diethylamine are added. The mixture is stirred at room temperature for 18 hours, then the solvent is removed *in vacuo*. The residue is purified by preparative HPLC.

10


Yield: 15 mg (75%)

<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ = 0.6 (t, 6H); 1.1 (t, 3H); 2.0 (m, 2H); 2.2 (m, 2H); 3.1 (br d, 1H); 3.9 (br d, 1H); 4.1 (q, 2H); 5.4 (d, 1H); 7.5 (m, 1H); 7.6 (m, 4H); 7.7 (m, 1H); 7.9 (m, 2H) ppm.

15

**Example 21**

Ethyl 6-(anilinomethyl)-4-(4-cyanophenyl)-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

50 mg (0.10 mmol) of Example 19 are dissolved in 2 ml acetone, and 18 mg (0.20 mmol) aniline are added. The mixture is stirred at room temperature for 18 hours, then the solvent is removed *in vacuo*. The residue is purified by preparative HPLC.

10


Yield: 28 mg (55%)

<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ = 1.1 (t, 3H); 3.6 (d/d, 1H); 4.1 (q, 2H); 4.4 (d/d, 1H); 5.4 (m, 2H); 6.2 (m, 2H); 6.5 (m, 1H); 6.9 (m, 2H); 7.6 (m, 6H); 7.9 (m, 2H); 8.4 (d, 1H) ppm.

15

**Example 22**

(+)-Ethyl 4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

The enantiomers of Example 1 are separated by preparative HPLC on a chiral phase: 100 mg compound dissolved in 1.5 ml ethyl acetate, column KBD 8361 (chiral silica gel selector based on monomer *N*-methacryloyl-L-leucine-1-menthylamide, cf. EP-A-379 917), 250 mm x 20 mm, eluent ethyl acetate, flow 25 ml/min, temperature 23°C, injection volume 2500 µl, detection 254 nm.

10

<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ = 1.1 (t, 3H); 2.0 (s, 3H); 4.0 (q, 2H); 5.4 (d, 1H); 7.6 (m, 3H); 7.7 (m, 2H); 7.8 (m, 1H); 7.9 (m, 2H); 8.4 (d, 1H) ppm.


[α]<sup>20</sup> = +3.3° (λ = 589 nm, dichloromethane, c = 535.0 mg / 100 ml)

15



**Example 23**

(-)-Ethyl 4-(4-cyanophenyl)-3,6-dimethyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

100 mg (0.23 mmol) of Example 22 are added to a suspension of 14 mg (0.35 mmol) 60% sodium hydride (in mineral oil) in 2 ml THF. The mixture is stirred at room temperature for two hours. Then 29 mg (0.23 mmol) dimethylsulfate are added, and the mixture is stirred at room temperature for another 2 hours. Then water and ethyl acetate are added, the organic phase is washed with water and brine, dried over sodium sulfate and evaporated *in vacuo*. The product is purified by column chromatography on silica with cyclohexane/ethyl acetate as eluent.

Yield: 76 mg (74%)


<sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>): δ = 1.1 (t, 3H); 2.0 (s, 3H); 2.8 (s, 3H); 4.0 (q, 2H); 5.5 (s, 1H); 7.6 (m, 3H); 7.7 (m, 1H); 7.8 (m, 2H); 7.9 (m, 2H) ppm.

[α]<sub>D</sub><sup>20</sup> = -18.1° (λ = 589 nm, dichloromethane, c = 530.0 mg / 100 ml)

15

**Example 24**

Ethyl 4-(6-cyano-3-pyridinyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

To a stirred solution of Example 3A (76 mg, 0.58 mmol) in tetrahydrofuran (5 ml) is given ethyl 3-oxobutanoate (75 mg, 0.58 mmol), N-[3-(trifluoromethyl)phenyl]urea (118 mg, 0.58 mmol) and polyphosphoric acid ethyl ester (200 mg; freshly prepared according to the procedure of Cava *et al.*, J. Org. Chem. 1969, 34, 2665). The reaction mixture is refluxed for two days (48 hours) after which time the solution is diluted with DMSO (2 ml) and purified by preparative HPLC. The product fractions are concentrated *in vacuo* and chromatographed again over silica with cyclohexane and ethyl acetate as eluent.

Yield: 92 mg (35% of th.)

MS (ESIpos):  $m/z = 431$  ( $M+H$ )<sup>+</sup>


HPLC (method 4) = 4.63 min

<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>):  $\delta$  = 8.76 (s, 1H), 8.36 (d, 1H), 8.16-8.00 (m, 2H), 7.83-7.74 (m, 2H), 7.75-7.58 (m, 2H), 5.47 (d, 1H), 4.03 (quartet, 2H), 2.06 (s, 3H), 1.08 (t, 3H) ppm.

20

**Example 25**

4-{5-(1H-Imidazol-1-ylcarbonyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-4-pyrimidinyl} benzonitrile



5

To a solution of 501 mg (1.25 mmol) of the compound of Example 11 in 5 ml dry dimethylformamide are added 567 mg (3.5 mmol) N,N-carbonyldiimidazole. After allowing the reaction mixture to stand overnight, the solvent is evaporated off *in vacuo*. The residue is taken up in ethyl acetate and washed with water and brine. After drying with magnesium sulfate the solvent is evaporated off *in vacuo*.

10

Yield: 500 mg (88.6% of th.)


MS (EI):  $m/z = 452$  (M+H)<sup>+</sup>

15

<sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>):  $\delta = 1.40$  (d, 3H), 5.5 (d, 1H), 7.0 (s, 1H), 7.55-8.0 (m, 9H), 8.4 (s, 1H), 8.45 (d, 1H) ppm.

**Example 26**

2-Hydroxyethyl 4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

45.1 mg (0.1 mmol) of the compound of Example 25 are added to 0.5 ml ethylene glycol. The reaction mixture is stirred at approx. 100°C for 1 hour. After cooling the reaction mixture is purified by preparative HPLC (column: Agilent Zorbax Extend C18 20 mm x 50 mm, 5 µm; solvent A: acetonitrile, solvent B: water + 0.1% conc. ammonia; gradient: 0 min 10% A, 2 min 10% A, 6 min 90% A, 7 min 90% A, 7.1 min 10% A, 8 min 10% A; wavelength: 220 nm; injection volume: approx. 500 µl; number of injections: 1). The product containing fractions are combined and concentrated *in vacuo*.

15 Yield: 22 mg (49.4% of th.)


MS (EI):  $m/z = 446$  ( $M+H$ )<sup>+</sup>

<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ = 2.05 (d, 3H), 3.5 (quartet, 2H), 3.95-4.15 (m, 2H), 4.75 (tr, 1H), 5.45 (d, 1H), 7.55-7.75 (m, 5H), 7.75 (d, 1H), 7.85 (d, 2H), 8.35 (d, 1H) ppm.

20

**Example 27**

2-(Dimethylamino)ethyl 4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)-phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

45.1 mg (0.1 mmol) of the compound of Example 25 are added to 0.5 ml 2-(dimethylamino)ethanol. The reaction mixture is stirred at approx. 100°C for 1 hour. After cooling the reaction mixture is purified by preparative HPLC (column: Agilent Zorbax Extend C18 20 mm x 50 mm, 5 µm; solvent A: acetonitrile, solvent B: water + 0.1% conc. ammonia; gradient: 0 min 10% A, 2 min 10% A, 6 min 90% A, 7 min 90% A, 7.1 min 10% A, 8 min 10% A; wavelength: 220 nm; injection volume: approx. 500 µl; number of injections: 1). The product containing fractions are combined and concentrated *in vacuo*.

15 Yield: 24 mg (50.8% of th.)


MS (EI):  $m/z = 473$  ( $M+H$ )<sup>+</sup>

<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ = 2.05 (d, 3H), 2.1 (s, 6H), 2.4 (m, 2H), 4.1 (m, 2H), 5.35 (d, 1H), 7.55 (d, 1H), 7.6 (d, 2H), 7.7 (m, 2H), 7.8 (d, 1H), 7.85 (d, 2H), 8.35 (d, 1H) ppm.

20

**Example 28**

2-(4-Pyridinyl)ethyl 4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate.



5

45.1 mg (0.1 mmol) of the compound of Example 25 are added to 0.5 ml 2-(4-pyridinyl)ethanol. The reaction mixture is stirred at approx. 100°C for 1 hour. After cooling the reaction mixture is purified by preparative HPLC (column: Agilent Zorbax Extend C18 20 mm x 50 mm, 5 µm; solvent A: acetonitrile, solvent B: water + 0.1% conc. ammonia; gradient: 0 min 10% A, 2 min 10% A, 6 min 90% A, 7 min 90% A, 7.1 min 10% A, 8 min 10% A; wavelength: 220 nm; injection volume: approx. 500 µl; number of injections: 1). The product containing fractions are combined and concentrated *in vacuo*.

15 Yield: 17 mg (33.5% of th.)


MS (EI):  $m/z = 507$  (M+H)<sup>+</sup>

<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ = 2.0 (d, 3H), 2.9 (tr, 2H), 4.3 (tr, 2H), 5.25 (d, 1H), 7.15 (d, 2H), 7.45 (d, 2H), 7.5 (d, 1H), 7.65 (tr, 2H), 7.8 (m, 3H), 8.35 (d, 1H), 8.4 (d, 2H) ppm.

20

**Example 29**

2-(2-Pyridinyl)ethyl 4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

45.1 mg (0.1 mmol) of the compound of Example 25 are added to 0.5 ml 2-(2-pyridinyl)ethanol. The reaction mixture is stirred at approx. 100°C for 1 hour. After cooling the reaction mixture is purified by preparative HPLC (column: Agilent Zorbax Extend C18 20 mm x 50 mm, 5 µm; solvent A: acetonitrile, solvent B: water + 0.1% conc. ammonia; gradient: 0 min 10% A, 2 min 10% A, 6 min 90% A, 7 min 90% A, 7.1 min 10% A, 8 min 10% A; wavelength: 220 nm; injection volume: approx. 500 µl; number of injections: 1). The product containing fractions are combined and concentrated *in vacuo*.

15 Yield: 22 mg (43.4% of th.)


MS (EI): m/z = 507 (M+H)<sup>+</sup>

<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ = 2.0 (d, 3H), 3.0 (tr, 2H), 4.4 (tr, 2H), 5.25 (d, 1H), 7.15-7.25 (m, 2H), 7.4 (d, 2H), 7.5 (d, 1H), 7.6-7.75 (m, 3H), 7.8 (m, 3H), 8.3 (d, 1H), 8.45 (d, 1H) ppm.

20

**Example 30**

2-(2-Oxo-1-pyrrolidinyl)ethyl 4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

45.1 mg (0.1 mmol) of the compound of Example 25 are added to 0.5 ml 1-(2-hydroxyethyl)-2-pyrrolidinone. The reaction mixture is stirred at approx. 100°C for 1 hour. After cooling the reaction mixture is purified by preparative HPLC (column: Agilent Zorbax Extend C18 20 mm x 50 mm, 5 µm; solvent A: acetonitrile, solvent B: water + 0.1% conc. ammonia; gradient: 0 min 10% A, 2 min 10% A, 6 min 90% A, 7 min 90% A, 7.1 min 10% A, 8 min 10% A; wavelength: 220 nm; injection volume: approx. 500 µl; number of injections: 1). The product containing fractions are combined and concentrated *in vacuo*.

10

Yield: 25 mg (48.8% of th.)





MS (EI):  $m/z = 513$  (M+H)<sup>+</sup>

<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ = 1.8 (quintet, 2H), 2.0 (d, 3H), 2.1 (tr, 2H), 3.2 (tr, 2H), 3.4 (tr, 2H), 4.0-4.2 (m, 2H), 5.35 (d, 1H), 7.55 (d, 1H), 7.6 (d, 2H), 7.7 (tr, 2H), 7.8 (d, 1H), 7.9 (d, 2H), 8.4 (d, 1H) ppm.





20

In analogy to the procedures for Examples 14-16, the following compounds are prepared:







Example No.	Structure	Starting materials	Yield [%]	R <sub>t</sub> [min] (method)	Mass [M+H] <sup>+</sup>
31		Example 1; ethyl bromoacetate	85	4.01 (1)	516
32		Example 1; cyclopropane- carbonyl chloride	79	4.09 (1)	498
33		Example 1; bromoethane	15	4.28 (2)	458
34		Example 1; 4-morpholine- carbonyl chloride	97	3.97 (2)	543

Example No.	Structure	Starting materials	Yield [%]	R <sub>t</sub> [min] (method)	Mass [M+H] <sup>+</sup>
35		Example 1; dimethylcarbamoyl chloride	98	4.00 (2)	523 [M+Na] <sup>+</sup>
36		Example 1; methyl chloridocarbonate	96	4.10 (2)	488
37		Example 1; benzylbromide	58	4.59 (2)	520
38		Example 1; propanoyl chloride	43	4.42 (2)	486




Example No.	Structure	Starting materials	Yield [%]	R <sub>t</sub> [min] (method)	Mass [M+H] <sup>+</sup>
39		Example 1; 2-methoxyethyl chlorido- carbonate	95	4.12 (2)	532
40		Example 1; isopropyl chlorido- carbonate	67	4.55 (2)	500
41		Example 1; diethylcarbamic chloride	18	4.25 (2)	529
42		Example 1; methyl (methyl- sulfonyl)- carbamic chloride	40	4.10 (2)	565



Example No.	Structure	Starting materials	Yield [%]	R <sub>t</sub> [min] (method)	Mass [M+H] <sup>+</sup>
43		Example 1; 2-bromo- acetamide; 2.5 equiv. NaH	54	3.7 (3)	487
44		Example 1; 2-bromoacetic acid; 2.5 equiv. NaH	67	3.8 (3)	488
45		Example 1; 2-bromo- ethanamine hydrobromide; 2.5 equiv. NaH	28	2.9 (2)	473
46		Example 1; 2-(chloro- methyl)pyridine hydrochloride; 2.5 equiv. NaH	37	4.0 (3)	521

Example No.	Structure	Starting materials	Yield [%]	R <sub>t</sub> [min] (method)	Mass [M+H] <sup>+</sup>
47		Example 1; N-(2-bromoethyl)-N,N-diethylamine hydrobromide; 2.5 equiv. NaH	82	2.98 (2)	529
48		Example 1; 2-bromo-N-methylacetamide; 2.5 equiv. NaH	65	3.70 (2)	501
49		Example 1; 3-(chloromethyl)pyridine hydrochloride; 2.5 equiv. NaH	15	3.68 (2)	521
50		Example 1; 4-(chloromethyl)pyridine hydrochloride; 2.5 equiv. NaH	21	3.47 (2)	521

Example No.	Structure	Starting materials	Yield [%]	R <sub>t</sub> [min] (method)	Mass [M+H] <sup>+</sup>
51		Example 1; 2-(bromo- methyl)-1H- imidazole hydrobromide; 2.5 equiv. NaH	6	2.97 (2)	510
52		Example 1; 3-(chloro- methyl)-1,2,4- oxadiazole	37	4.0 (3)	469
53		Example 1; 2-bromo-N-(2- methoxyethyl)- acetamide	91	3.77 (2)	545


In analogy to the procedures for Examples 6-8, the following compounds are prepared:

Example No.	Structure	Starting materials	Yield [%]	R <sub>t</sub> [min] (method)	Mass [M+H] <sup>+</sup>
54		N-[3-(trifluoro- methyl)phenyl]-urea; 4-cyano- benzaldehyde; methyl 3-oxobutanoate	79	3.68 (2)	416
55		N-[3-(trifluoro- methyl)phenyl]-urea; 4-cyano- benzaldehyde; cyclopropylmethyl 3-oxobutanoate	58	4.09 (2)	456
56		N-[3-(trifluoro- methyl)phenyl]-urea; 4-cyano- benzaldehyde; isopropyl 3-oxobutanoate	85	4.03 (2)	444

Example No.	Structure	Starting materials	Yield [%]	R <sub>t</sub> [min] (method)	Mass [M+H] <sup>+</sup>
57		N-[3-(trifluoromethyl)phenyl]-urea; 4-cyano-benzaldehyde; (1R)-2-methoxy-1-methyl-2-oxo-ethyl 3-oxobutanoate	73	3.82 (2)	488
58		N-[3-(trifluoromethyl)phenyl]-urea; 4-cyano-benzaldehyde; N,N-dimethyl-3-oxobutanamide	9	3.22 (2)	429

**Example 59**

5 Ethyl 4-(4-cyanophenyl)-6-methyl-3-[2-(4-morpholinyl)-2-oxoethyl]-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate








80 mg (0.16 mmol) of Example 44 are dissolved in 2 ml dimethylformamide, 16 mg (0.18 mmol) morpholine, 24 mg (0.18 mmol) 1-hydroxy-1H-benzotriazole hydrate and 20 mg (0.16 mmol) 4-dimethylaminopyridine are added. The reaction mixture is stirred at 0°C, then 35 mg (0.18 mmol) 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride are added. The reaction mixture is stirred at room temperature for 18 hours, then water and ethyl acetate are added. The organic phase is dried over sodium sulfate and evaporated to dryness *in vacuo*. If necessary, the product is further purified by column chromatography or preparative HPLC.





Yield: 78 mg (85%)




<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>):  $\delta$  = 1.1 (t, 3H); 2.0 (s, 3H); 3.4 (m, 4H); 3.6 (m, 4H); 3.7 (d, 1H); 4.1 (m, 2H); 4.5 (d, 1H); 5.5 (s, 1H); 7.6 (m, 5H); 7.8 (m, 1H); 7.9 (m, 2H) ppm.

In analogy to the procedure for Example 59, the following compounds are prepared:

Example No.	Structure	Starting materials	Yield [%]	R <sub>t</sub> [min] (method)	Mass [M+H] <sup>+</sup>
60		Example 44; N-methyl-piperazine	90	2.93 (2)	570
61		Example 44; N-(2-aminoethyl)-N,N-dimethyl-amine	87	2.93 (2)	558
62		Example 44; dimethyl-amine (2 M in THF)	83	3.84 (2)	515


In analogy to the procedures for Examples 6-8, the following compounds are  
5 prepared:

Example No.	Structure	Starting materials	Yield [%]	R <sub>f</sub> [min] (method)	Mass [M+H] <sup>+</sup>
63		N-[3-(trifluoromethyl)phenyl]-urea; 4-cyano-benzaldehyde; 1-(3-methyl-1,2,4-oxadiazol-5-yl)acetone	23	3.80 (3)	440
64		N-[3-(trifluoromethyl)phenyl]-urea; 4-cyano-benzaldehyde; 1-(1,3-benzothiazol-2-yl)-acetone	23	4.42 (2)	491
65		N-[3-(trifluoromethyl)phenyl]-urea; 4-cyano-benzaldehyde; 5-methyl-2,4-hexanedione	33	4.3 (1)	428
66		N-[3-(trifluoromethyl)phenyl]-urea; 4-cyano-benzaldehyde; 1-methoxy-2,4-pentanedione	3	3.47 (2)	430

Example No.	Structure	Starting materials	Yield [%]	R <sub>t</sub> [min] (method)	Mass [M+H] <sup>+</sup>
67		N-[3-(trifluoromethyl)phenyl]-urea; 4-cyano-benzaldehyde; 1-(2-furyl)-1,3-butanedione	13	3.70 (2)	452
68		N-[3-(trifluoromethyl)phenyl]-urea; 4-cyano-benzaldehyde; 1-phenyl-1,3-butanedione	14	4.03 (2)	462
69		N-[3-(trifluoromethyl)phenyl]-urea; 4-cyano-benzaldehyde; 1,1,1-trifluoro-2,4-pentanedione	5	3.9 (3)	454

**Example 70**

4-(4-Cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxamide



5

200 mg (0.5 mmol) of Example 11 are dissolved in 5 ml tetrahydrofuran and 6 mg (0.05 mmol) 4-N,N-dimethylaminopyridine, 77 mg (0.6 mmol) N,N-diisopropylethylamine and 115 mg (0.6 mmol) benzotriazol-1-yloxy-tris(pyrrolidino)phosphonium hexafluorophosphate are added. The reaction mixture is stirred at room temperature for 15 minutes, then 5 ml (2.5 mmol) ammonia (as 0.5 M solution in dioxane) are added. The reaction mixture is stirred at room temperature for 1 hour, then water and ethyl acetate are added. The organic phase is dried over sodium sulfate and evaporated to dryness *in vacuo*. The product is further purified by preparative HPLC.


15

Yield: 55 mg (28% of th.)

<sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>): δ = 1.8 (s, 3H); 5.4 (d, 1H); 7.2 (br. s, 1H); 7.4 (br. s, 1H); 7.6 (m, 5H); 7.7 (m, 1H); 7.9 (m, 2H); 8.1 (d, 1H) ppm.

**Example 71**

(+)-4-(4-Cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylic acid



5


The enantiomers of Example 11 are separated by preparative HPLC on a chiral phase [column KBD 8361 (chiral silica gel selector based on monomer *N*-methacryloyl-L-leucine-1-menthylamide, cf. EP-A-379 917), 250 mm x 20 mm, eluent: ethyl acetate  
 10 → methanol → ethyl acetate, flow 25 ml/min, temperature 23°C, detection 254 nm].

<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ = 2.0 (s, 3H); 5.4 (d, 1H); 7.6 (m, 1H); 7.6 (m, 2H); 7.7 (m, 1H); 7.8 (m, 1H); 7.9 (m, 3H); 8.3 (d, 1H); 12.5 (s, 1H) ppm.

[α]<sup>20</sup> = +2.5° (λ = 589 nm, methanol, c = 505 mg / 100 ml).

**Example 72**

(+)-2-Hydroxyethyl 4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)-phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

Under argon, 1560 mg (3.89 mmol) of the compound of Example 71 are added to 19.6 ml DMF. After addition of 1.095 ml (7.86 mmol) triethylamine and 1.11 ml (15.7 mmol) 2-bromoethanol, the reaction mixture is stirred at ca. 70°C for 8 hours.

10 After cooling, the reaction mixture is concentrated *in vacuo*. The residue is taken up in ethyl acetate and washed with water. After drying with magnesium sulfate, the organic phase is evaporated *in vacuo*. The residue is taken up in 8 ml methanol and purified by preparative HPLC (column: Nucleosil 100-5 C 18 Nautilus, 20 x 50 mm, 5 µm; solvent A: acetonitrile, solvent B: water + 0.3% formic acid; gradient: 0 min 10% A, 2 min 10% A, 6 min 90% A, 7 min 90% A, 7.1 min 10% A, 8 min 10% A; wavelength: 220 nm; injection volume: ca. 500 µl; number of injections: 18). The product containing fractions are combined and lyophilized.

15

Yield: 1290 mg (74.5% of th.)


MS (EI):  $m/z = 446 (M+H)^+$

20 <sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ = 2.05 (d, 3H); 3.5 (quartett, 2H); 3.95-4.15 (m, 2H); 4.75 (tr, 1H); 5.45 (d, 1H); 7.55-7.75 (m, 5H); 7.75 (d, 1H); 7.85 (d, 2H); 8.35 (d, 1H) ppm.

[α]<sup>20</sup> = +14.3° (λ = 589 nm, methanol, c = 455 mg / 100 ml).

**Example 73**

5-{5-Acetyl-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-4-pyrimidinyl}-2-pyridinecarbonitrile



5

To a stirred solution of Example 3A (75 mg, 0.57 mmol) in tetrahydrofuran (5 ml) is given 2,4-pentandione (57 mg, 0.57 mmol), N-[3-(trifluoromethyl)phenyl]urea (116 mg, 0.57 mmol) and polyphosphoric acid ethyl ester (200 mg) [freshly prepared according to the procedure of Cava *et al.*, J. Org. Chem. 34, 2665 (1969)]. The reaction mixture is refluxed for 24 hours after which time the solution is diluted with DMSO (2 ml) and purified by preparative HPLC.

10

Yield: 101 mg (44% of th.)


15

<sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>): δ = 2.02 (s, 3H); 2.24 (s, 3H); 5.54 (d, 1H); 7.52-7.90 (m, 4H); 8.08 (d, 2H); 8.50 (d, 1H); 8.81 (s, 1H) ppm.



**Example 74**

(+)-5-{5-Acetyl-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-4-pyrimidinyl}-2-pyridinecarbonitrile



5

The enantiomers of Example 73 are separated by preparative HPLC on a chiral phase [column KBD 8361 (chiral silica gel selector based on monomer *N*-methacryloyl-L-leucine-1-menthylamide, cf. EP-A-379 917), 250 mm x 20 mm, eluent: ethyl acetate → methanol → ethyl acetate, flow 25 ml/min, temperature 23°C, detection 254 nm].

10

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>): δ = 2.06 (s, 3H); 2.35 (s, 3H); 5.69 (d, 1H); 6.02 (d, 1H); 7.29-7.50 (m, 2H); 7.57-7.75 (m, 3H); 7.83 (dd, 1H); 8.74 (d, 1H) ppm.


MS (ESIpos): m/z = 401 (M+H)<sup>+</sup>

[α]<sub>D</sub><sup>20</sup> = +25.1° (δ = 589 nm, methanol, c = 505 mg / 100 ml).

15

**Example 75**

2-(2-Pyridinyl)methyl 4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)-phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



To a solution of 40.1 mg (0.1 mmol) of the compound of Example 11 in 0.4 ml dry dimethylformamide are added 48.6 mg (0.3 mmol) N,N-carbonyldiimidazole. After allowing the reaction mixture to stand for one hour, the reaction mixture is diluted with water and extracted with dichloromethane. After drying with magnesium sulfate, the solvent is evaporated off *in vacuo*. To the residue are added to 0.5 ml (2-pyridinyl)methanol. The reaction mixture is stirred at approx. 100°C for 1 hour. After cooling, the reaction mixture is purified by preparative HPLC (column: Nucleosil 100-5 C 18 Nautilus 20 mm x 50 mm, 5 µm; solvent A: acetonitrile, solvent B: water + 0.1% formic acid; gradient: 0 min 10% A, 2 min 10% A, 6 min 90% A, 7 min 90% A, 7.1 min 10% A, 8 min 10% A; flow rate 25 ml/min; wavelength: 220 nm; injection volume: approx. 550 µl; number of injections: 1). The product containing fractions are combined and concentrated *in vacuo*.


Yield: 17 mg (34.5% of th.)

MS (EI):  $m/z = 493$  (M+H)<sup>+</sup>

<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ = 2.1 (d, 3H); 5.15 (dd, 2H); 5.45 (d, 1H); 7.05 (d, 1H); 7.3 (dd, 1H); 7.5-7.85 (m, 9H); 8.35 (d, 1H); 8.5 (d, 2H) ppm.

**Example 76**

2-(3-Pyridinyl)ethyl 4-(4-cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)-phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

To a solution of 60.2 mg (0.15 mmol) of the compound of Example 11 in 0.57 ml dry dimethylformamide are added 72.9 mg (0.45 mmol) N,N-carbonyldiimidazole. After allowing the reaction mixture to stand for one hour, the reaction mixture is diluted with water and extracted with ethylacetate. After drying with magnesium sulfate, the solvent is evaporated off *in vacuo*. To the residue are added 185 mg (1.5 mmol) 2-(3-pyridyl)ethanol and 20  $\mu$ l (0.27 mmol) triethylamine. The reaction mixture is stirred for one hour at 100°C. Then the reaction mixture is diluted with 0.4 ml methanol, filtered and purified by preparative HPLC (column: Nucleosil 100-5 C 18 Nautilus


10 20 mm x 50 mm, 5  $\mu$ m; solvent A: acetonitrile, solvent B: water + 0.1% formic acid; gradient: 0 min 10% A, 2 min 10% A, 6 min 90% A, 7 min 90% A, 7.1 min 10% A, 8 min 10% A; flow rate 25 ml/min; wavelength: 220 nm; injection volume: approx. 550  $\mu$ l; number of injections: 1). The product containing fractions are combined and concentrated *in vacuo*.

20 Yield: 44 mg (57.9% of th.)

LC-MS (EI, method 5):  $m/z$  = 507 (M+H)<sup>+</sup>,  $R_t$  = 3.19 min.

**Example 77**

4-(4-Cyanophenyl)-3,6-dimethyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylic acid



5

4.1 g (9.25 mmol) of Example 14 are dissolved in 100 ml ethanol. To this solution 6.2 ml (27.6 mmol) of a solution of potassium hydroxide in water (25% by weight) are added. The reaction mixture is allowed to stand at room temperature for 18 hours.

10 Then further 12.4 ml (55.2 mmol) of a solution of potassium hydroxide in water (25% by weight) are added and the reaction mixture is stirred for 2 hours. The reaction mixture is diluted with water and extracted three times with ethyl acetate. The aqueous phase is acidified with 1 N hydrochloric acid and extracted with ethyl acetate. This last extract is dried over magnesium sulfate and evaporated off *in vacuo*. The residue is purified by column chromatography on silica with cyclohexane/ethyl acetate as eluent.

15

Yield: 1.5 g (39% of th.)





MS (EI):  $m/z = 416$  ( $M+H$ )<sup>+</sup>





<sup>1</sup>H-NMR (300 MHz, DMSO- $d_6$ ):  $\delta = 2.0$  (s, 3H); 2.8 (s, 3H); 5.5 (d, 1H); 7.6-7.8 (m, 6H); 7.9(d, 2H); 12.6 (s, 1H) ppm.

20

In analogy to the procedure for Example 76, the following compounds are prepared:





Example No.	Structure	Starting materials	Yield [%]	R <sub>t</sub> [min] (method)	Mass [M+H] <sup>+</sup>
78		Example 11; 3-pyridinyl-methanol	56.9	3.45 (5)	493
79		Example 11; 2-hydroxy-acetamide <sup>1)</sup>	61.1	3.38 (5)	459
80		Example 11; 2-hydroxyethyl-(methyl)formamide	80.9	3.5 (5)	487
81		Example 11; 2-hydroxyethyl-acetamide	56.2	3.44 (5)	487





Example No.	Structure	Starting materials	Yield [%]	R <sub>t</sub> [min] (method)	Mass [M+H] <sup>+</sup>
82		Example 11; (1-methyl-1H-imidazol-5-yl)-methanol <sup>1)</sup>	45.8	2.87 (5)	496
83		Example 11; 2-(1H-pyrazol-1-yl)ethanol	60.6	3.7 (5)	496
84		Example 11; 2-(1H-1,2,4-triazol-1-yl)-ethanol <sup>1)</sup>	67.1	3.48 (5)	497
85		Example 11; 2-hydroxyethyl acetate	56.1	3.98 (5)	488

Example No.	Structure	Starting materials	Yield [%]	R <sub>t</sub> [min] (method)	Mass [M+H] <sup>+</sup>
86		Example 11; 2-(dimethyl- amino)-2- methyl-1- propanol	34.6	2.9 (5)	502
87		Example 11; 3-(dimethyl- amino)propanol	54.8	2.86 (5)	487
88		Example 11; 2-(1- pyrrolidinyl)- ethanol	56.2	2.86 (5)	500
89		Example 77; 2-(3-pyridinyl)- ethanol	58.9	3.36 (5)	522

Example No.	Structure	Starting materials	Yield [%]	R <sub>t</sub> [min] (method)	Mass [M+H] <sup>+</sup>
90		Example 77; (3-pyridinyl)- methanol	61.9	3.64 (5)	507
91		Example 77; 2-hydroxy- acetamide <sup>1)</sup>	53.6	3.54 (5)	473
92		Example 77; 2-hydroxyethyl- (methyl)form- amide	54.6	3.68 (5)	501
93		Example 77; 2-hydroxyethyl- acetamide	66.6	3.59 (5)	501




Example No.	Structure	Starting materials	Yield [%]	R <sub>t</sub> [min] (method)	Mass [M+H] <sup>+</sup>
94	 2	Example 77; (1-methyl-1H-imidazol-5-yl)-methanol <sup>1)</sup>	34.0	3.02 (5)	510
95		Example 77; 2-(1H-pyrazol-1-yl)ethanol	61.5	3.91 (5)	510
96	 E24	Example 77; 2-(1H-1,2,4-triazol-1-yl)-ethanol <sup>1)</sup>	71.8	3.64 (5)	511
97	 5	Example 77; 2-hydroxyethyl acetate	53.2	4.12 (5)	502

Example No.	Structure	Starting materials	Yield [%]	R <sub>t</sub> [min] (method)	Mass [M+H] <sup>+</sup>
98		Example 77; 2-(dimethyl- amino)-2- methyl-1- propanol	25.9	3.02 (5)	516
99		Example 77; 3-(dimethyl- amino)propanol	54.6	2.98 (5)	502
100		Example 77; 2-(1- pyrrolidinyl)- ethanol	55.9	2.98 (5)	514
101		Example 77; (2-pyridinyl)- methanol	67.1	3.91 (5)	507

<sup>1)</sup> in this case the alcohol used is a solid and the reaction is conducted in the presence of 0.4 ml DMF

**Example 102**

Ethyl 4-(4-cyanophenyl)-1-(3,5-dichlorophenyl)-6-methyl-2-oxo-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate



5

Under argon, 30.8 mg (0.15 mmol) N-(3,5-dichlorophenyl)urea are stirred together with 39.3 mg (0.3 mmol) 4-formylbenzonitrile, 39 mg (0.3 mmol) ethyl 3-oxo-butanoate and 90 mg trimethylsilylpolyphosphate in 0.5 ml dioxan at 80°C for 4 hours. After adding a small amount of DMSO, the reaction mixture is filtered and purified by preparative HPLC (column: Agilent Zorbax Extend C18 20 mm x 50 mm, 5 µm; solvent A: acetonitrile, solvent B: water + 0.1% conc. aq. ammonia; gradient: 0 min 10% A, 2 min 10% A, 6 min 90% A, 7 min 90% A, 7.1 min 10% A, 8 min 10% A; flow rate 25 ml/min; wavelength: 220 nm; injection volume: approx. 500 µl; number of injections: 1). The product containing fractions are combined and concentrated *in vacuo*.


15

Yield: 38.1 mg (59% of th.)

LC-MS (EI, method 7):  $m/z = 431$  ( $M+H$ )<sup>+</sup>,  $R_t = 4.14$  min.

**Example 103**

Ethyl 6-methyl-4-(3-nitrophenyl)-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate







30.6 mg (0.15 mmol) N-[3-(trifluoromethyl)phenyl]urea are shaken together with 45.3 mg (0.3 mmol) 3-nitrobenzaldehyde, 39 mg (0.3 mmol) ethyl 3-oxobutanoate and 90 mg polyphosphoric acid ethyl ester [freshly prepared according to the procedure of Cava *et al.*, J. Org. Chem. 34, 2665 (1969)] in 0.5 ml dioxan and 0.1 ml DMF at 80°C for 18 hours. After adding 200 µl DMF, the reaction mixture is filtered and purified by preparative HPLC (column: Nucleosil 100-5 C 18 Nautilus 20 mm x 50 mm, 5 µm; solvent A: acetonitrile, solvent B: water + 0.1% formic acid; gradient: 0 min 10% A, 2 min 10% A, 6 min 90% A, 7 min 90% A, 7.1 min 10% A, 8 min 10% A; flow rate 25 ml/min; wavelength: 220 nm; injection volume: approx. 800 µl; number of injections: 1). The product containing fractions are combined and concentrated *in vacuo*.

Yield: 34 mg (50.4% of th.)


LC-MS (EI, method 6):  $m/z = 450 (M+H)^+$ ,  $R_t = 3.94$  min.

In analogy to the procedure for Example 102, the following compounds are prepared:

Example No.	Structure	Starting materials	Yield [%]	R <sub>t</sub> [min] (method)	Mass [M+H] <sup>+</sup>
104		N-(3-nitro-phenyl)urea; 4-chlorobenzaldehyde; ethyl 3-oxobutanoate	70.5	3.65 (6)	417
105		N-(3-nitro-phenyl)urea; 3-nitrobenzaldehyde; ethyl 3-oxobutanoate	81.3	3.61 (6)	427
106		N-(3-nitro-phenyl)urea; 4-fluorobenzaldehyde; ethyl 3-oxobutanoate	56.8	3.63 (6)	400
107		N-(3-nitro-phenyl)urea; 4-bromobenzaldehyde; ethyl 3-oxobutanoate	69.5	4.02 (5)	461

**Example 108**

4-(4-Cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydro-pyrimidine-5-carboxylic acid 2-cyanoethyl ester



5

9.87 g (48.3 mmol) N-[3-(trifluoromethyl)phenyl]urea, 12.68 g (96.68 mmol) 4-cyanobenzaldehyde, 15 g (96.68 mmol) (2-cyanoethyl) 3-oxobutanoate and 37.5 g polyphosphoric acid ethyl ester are suspended in 250 ml of THF. The mixture is stirred at reflux for 18 hours. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with cyclohexane/ethyl acetate as eluent.

10


Yield: 25 g (100% of th.)

<sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>): δ = 2.1 (s, 3H); 2.8 (m, 2H); 4.2 (m, 2H); 5.4 (d, 1H); 7.6 (m, 4H); 7.7 (m, 2H); 7.9 (m, 2H); 8.5 (d, 1H) ppm.

15

**Example 109**

4-(4-Cyanophenyl)-6-methyl-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidine-5-carbonitrile



5

0.609 g (1.52 mmol) of Example 70 are dissolved in 60 ml THF and 1.24 g (12.93 mmol) (methoxycarbonylsulfamoyl)-triethylammonium-N-betaine are added. The reaction mixture is stirred at room temperature for 1 hour, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with dichloromethane/methanol mixtures as eluent.

10


Yield: 249 mg (43% of th.)

<sup>1</sup>H-NMR (300 MHz, DMSO-d<sub>6</sub>): δ = 1.8 (s, 3H); 5.4 (d, 1H); 7.7 (m, 4H); 7.8 (m, 2H); 8.0 (m, 2H), 8.4 (d, 1H) ppm.

15

**Example 110**

Ethyl 6-methyl-4-(4-nitrophenyl)-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidine-5-carboxylate



5

7.84 g (38.4 mmol) N-[3-(trifluoromethyl)phenyl]urea, 5.81 g (38.4 mmol) 4-nitrobenzaldehyde, 5.0 g (38.4 mmol) ethyl 3-oxobutanoate and 15 g polyphosphoric acid ethyl ester are suspended in 100 ml of THF. The mixture is stirred at reflux for 18 hours. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with toluene/ethyl acetate as eluent.

10

Yield: 8.75 g (51% of th.)


<sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>): δ = 1.1 (t, 3H); 2.1 (s, 3H); 4.0 (m, 2H); 5.4 (d, 1H); 7.5-7.8 (m, 6H); 8.3 (m, 2H); 8.5 (d, 1H) ppm.

15



**Example 111**

5-Acetyl-6-methyl-4-(4-nitrophenyl)-2-oxo-1-[3-(trifluoromethyl)phenyl]-1,2,3,4-tetrahydropyrimidine



5

0.407 g (2.0 mmol) N-[3-(trifluoromethyl)phenyl]urea, 0.302 g (2.0 mmol) 4-nitrobenzaldehyde, 0.2 g (2.0 mmol) 2,4-pentanedione and 0.4 g polyphosphoric acid ethyl ester are suspended in 20 ml of THF. The mixture is stirred at reflux for 18 hours. After cooling down to room temperature, the solvent is removed *in vacuo* and the residue is purified by column chromatography on silica with cyclohexane/ethyl acetate as eluent.

10

Yield: 0.302 g (36% of th.)

<sup>1</sup>H-NMR (200 MHz, DMSO-d<sub>6</sub>): δ = 2.0 (s, 3H); 2.2 (s, 3H); 5.5 (d, 1H); 7.5-7.8 (m, 6H); 8.3 (m, 2H); 8.5 (d, 1H) ppm.

15

### C. Operative examples relating to pharmaceutical compositions

5 The compounds according to the invention can be converted into pharmaceutical preparations as follows:

#### Tablet

##### Composition

10

100 mg of the compound of Example 1, 50 mg of lactose (monohydrate), 50 mg of maize starch (native), 10 mg of polyvinylpyrrolidone (PVP 25) (from BASF, Ludwigshafen, Germany) and 2 mg of magnesium stearate.

Tablet weight 212 mg, diameter 8 mm, curvature radius 12 mm.

15

##### Preparation

20 The mixture of active component, lactose and starch is granulated with a 5% solution (m/m) of the PVP in water. After drying, the granules are mixed with magnesium stearate for 5 min. This mixture is moulded using a customary tablet press (tablet format, see above). The moulding force applied is typically 15 kN.

#### Orally administrable suspension

25

##### Composition

1000 mg of the compound of Example 1, 1000 mg of ethanol (96%), 400 mg of Rhodigel (xanthan gum from FMC, Pennsylvania, USA) and 99 g of water.

30


A single dose of 100 mg of the compound according to the invention is provided by 10 ml of oral suspension.

Preparation

5 The Rhodigel is suspended in ethanol and the active component is added to the suspension. The water is added with stirring. Stirring is continued for about 6h until the swelling of the Rhodigel is complete.

**We claim**

1. Compounds of the general formula (I)



wherein

A represents an aryl or heteroaryl ring,

$R^1$ ,  $R^2$  and  $R^3$  independently from each other represent hydrogen, halogen, nitro, cyano,  $C_1$ - $C_6$ -alkyl, hydroxy or  $C_1$ - $C_6$ -alkoxy, wherein  $C_1$ - $C_6$ -alkyl and  $C_1$ - $C_6$ -alkoxy can be further substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy and  $C_1$ - $C_4$ -alkoxy,

$R^4$  represents trifluoromethylcarbonyl,  $C_1$ - $C_6$ -alkylcarbonyl,  $C_1$ - $C_6$ -alkoxycarbonyl,  $C_1$ - $C_6$ -alkenoxycarbonyl, hydroxycarbonyl, aminocarbonyl, mono- or di- $C_1$ - $C_4$ -alkylaminocarbonyl,  $C_6$ - $C_{10}$ -arylaminocarbonyl, arylcarbonyl, heteroarylcarbonyl, heterocyclylcarbonyl, heteroaryl, heterocyclyl or cyano, wherein  $C_1$ - $C_6$ -alkylcarbonyl,  $C_1$ - $C_6$ -alkoxycarbonyl, mono- and di- $C_1$ - $C_4$ -alkylaminocarbonyl can be further substituted with one to three identical or different radicals selected from the group consisting of  $C_3$ - $C_8$ -cycloalkyl, hydroxy,

C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl, hydroxycarbonyl, amino-carbonyl, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, C<sub>1</sub>-C<sub>4</sub>-alkyl-carbonylamino, (C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyl)-C<sub>1</sub>-C<sub>4</sub>-alkylamino, cyano, amino, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylamino, heteroaryl, heterocyclyl and tri-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-silyl, and wherein heteroarylcarbonyl, heterocyclyl-carbonyl, heteroaryl and heterocyclyl can be further substituted with C<sub>1</sub>-C<sub>4</sub>-alkyl,

R<sup>5</sup> represents C<sub>1</sub>-C<sub>4</sub>-alkyl, which can be substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-alkenoxo, C<sub>1</sub>-C<sub>6</sub>-alkylthio, amino, mono- and di-C<sub>1</sub>-C<sub>6</sub>-alkylamino, arylamino, hydroxycarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl and the radical -O-C<sub>1</sub>-C<sub>4</sub>-alkyl-O-C<sub>1</sub>-C<sub>4</sub>-alkyl,

or

R<sup>5</sup> represents amino,

R<sup>6</sup> represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, formyl, aminocarbonyl, mono- or di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, C<sub>3</sub>-C<sub>8</sub>-cycloalkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkyl-carbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, N-(C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl)-amino-carbonyl, N-(C<sub>1</sub>-C<sub>4</sub>-alkylsulfonyl)-N-(C<sub>1</sub>-C<sub>4</sub>-alkyl)-aminocarbonyl, heteroaryl, heterocyclyl, heteroarylcarbonyl or heterocyclylcarbonyl, wherein C<sub>1</sub>-C<sub>6</sub>-alkyl, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, heteroaryl and heterocyclyl can be substituted with one to three identical or different radicals selected from the group consisting of aryl, heteroaryl, hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, hydroxycarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, aminocarbonyl, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, amino, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkyl-amino, C<sub>1</sub>-C<sub>4</sub>-alkylcarbonylamino, tri-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-silyl, cyano,


- 105 -

mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylamino-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl and halogen,

or

5

R<sup>6</sup> represents a moiety of the formula



wherein

10

R<sup>6A</sup> is selected from the group consisting of hydrogen and C<sub>1</sub>-C<sub>6</sub>-alkyl, and

n represents an integer of 1 or 2,

15

R<sup>7</sup> represents halogen, nitro, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, hydroxy or C<sub>1</sub>-C<sub>6</sub>-alkoxy, wherein C<sub>1</sub>-C<sub>6</sub>-alkyl and C<sub>1</sub>-C<sub>6</sub>-alkoxy can be further substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy and C<sub>1</sub>-C<sub>4</sub>-alkoxy,

20

and

Y<sup>1</sup>, Y<sup>2</sup>, Y<sup>3</sup>, Y<sup>4</sup> and Y<sup>5</sup> independently from each other represent CH or N, wherein the ring contains either 0, 1 or 2 nitrogen atoms,

25

and their salts, hydrates and/or solvates and their tautomeric forms.

2. Compounds of general formula (I) according to Claim 1, wherein

A represents an aryl or heteroaryl ring,

5 R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> independently from each other represent hydrogen, halogen, nitro, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, hydroxy or C<sub>1</sub>-C<sub>6</sub>-alkoxy, wherein C<sub>1</sub>-C<sub>6</sub>-alkyl and C<sub>1</sub>-C<sub>6</sub>-alkoxy can be further substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy and C<sub>1</sub>-C<sub>4</sub>-alkoxy,

10 R<sup>4</sup> represents C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkenoxycarbonyl, hydroxycarbonyl, aminocarbonyl, mono- or di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, C<sub>6</sub>-C<sub>10</sub>-arylaminocarbonyl, heteroarylcarbonyl, heterocyclylcarbonyl, heteroaryl, heterocyclyl or cyano, wherein C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl can be further substituted with one to three  
15 identical or different radicals selected from the group consisting of C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl, hydroxycarbonyl, aminocarbonyl, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, C<sub>1</sub>-C<sub>4</sub>-alkylcarbonylamino, amino, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylamino, heteroaryl, heterocyclyl and tri-(C<sub>1</sub>-C<sub>6</sub>-alkyl)-silyl,  
20

R<sup>5</sup> represents C<sub>1</sub>-C<sub>4</sub>-alkyl, which can be substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy, C<sub>1</sub>-C<sub>6</sub>-alkoxy, C<sub>1</sub>-C<sub>6</sub>-alkenoxy, C<sub>1</sub>-C<sub>6</sub>-alkylthio,  
25 amino, mono- and di-C<sub>1</sub>-C<sub>6</sub>-alkylamino, arylamino, hydroxycarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl and the radical -O-C<sub>1</sub>-C<sub>4</sub>-alkyl-O-C<sub>1</sub>-C<sub>4</sub>-alkyl,


or

30 R<sup>5</sup> represents amino,

$R^6$  represents hydrogen,  $C_1$ - $C_6$ -alkyl, formyl, aminocarbonyl, mono- or di- $C_1$ - $C_4$ -alkylaminocarbonyl,  $C_3$ - $C_8$ -cycloalkylcarbonyl,  $C_1$ - $C_6$ -alkylcarbonyl,  $C_1$ - $C_6$ -alkoxycarbonyl, N-( $C_1$ - $C_4$ -alkylsulfonyl)-aminocarbonyl, N-( $C_1$ - $C_4$ -alkylsulfonyl)-N-( $C_1$ - $C_4$ -alkyl)-aminocarbonyl, heteroaryl, heterocyclyl, heteroarylcarbonyl or heterocyclylcarbonyl, wherein  $C_1$ - $C_6$ -alkyl, mono- and di- $C_1$ - $C_4$ -alkylaminocarbonyl,  $C_1$ - $C_6$ -alkylcarbonyl,  $C_1$ - $C_6$ -alkoxycarbonyl, heteroaryl and heterocyclyl can be substituted with one to three identical or different radicals selected from the group consisting of aryl, heteroaryl, hydroxy,  $C_1$ - $C_4$ -alkoxy, hydroxycarbonyl,  $C_1$ - $C_6$ -alkoxycarbonyl, aminocarbonyl, mono- and di- $C_1$ - $C_4$ -alkylaminocarbonyl, amino, mono- and di- $C_1$ - $C_4$ -alkylamino,  $C_1$ - $C_4$ -alkylcarbonylamino, tri-( $C_1$ - $C_6$ -alkyl)-silyl, cyano, mono- and di- $C_1$ - $C_4$ -alkylamino- $C_1$ - $C_4$ -alkylaminocarbonyl,  $C_1$ - $C_4$ -alkoxy- $C_1$ - $C_4$ -alkylaminocarbonyl and halogen,

or

$R^6$  represents a moiety of the formula



wherein

$R^{6A}$  is selected from the group consisting of hydrogen and  $C_1$ - $C_6$ -alkyl, and

$n$  represents an integer of 1 or 2,



R<sup>7</sup> represents halogen, nitro, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, hydroxy or C<sub>1</sub>-C<sub>6</sub>-alkoxy, wherein C<sub>1</sub>-C<sub>6</sub>-alkyl and C<sub>1</sub>-C<sub>6</sub>-alkoxy can be further substituted with one to three identical or different radicals selected from the group consisting of halogen, hydroxy and C<sub>1</sub>-C<sub>4</sub>-alkoxy,

5

and

Y<sup>1</sup>, Y<sup>2</sup>, Y<sup>3</sup>, Y<sup>4</sup> and Y<sup>5</sup> independently from each other represent CH or N, wherein the ring contains either 0, 1 or 2 nitrogen atoms.

10

3. Compounds of general formula (I) according to Claim 1 or 2, wherein

A represents a phenyl, naphthyl or pyridyl ring,

15

R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> independently from each other represent hydrogen, fluoro, chloro, bromo, nitro, cyano, methyl, ethyl, trifluoromethyl or trifluoromethoxy,

20

R<sup>4</sup> represents C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, hydroxycarbonyl, aminocarbonyl, mono-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl or cyano, wherein C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl and mono-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl can be substituted with one to three identical or different radicals selected from the group consisting of C<sub>3</sub>-C<sub>8</sub>-cycloalkyl, hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl, amino, mono- or di-C<sub>1</sub>-C<sub>4</sub>-alkylamino, heteroaryl and heterocyclyl,

25

R<sup>5</sup> represents methyl or ethyl,

30


R<sup>6</sup> represents hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, mono- or di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkylcarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl or heterocyclcarbonyl, wherein C<sub>1</sub>-C<sub>6</sub>-alkyl and C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl can be

- 109 -

substituted with one to three identical or different radicals selected from the group consisting of heteroaryl, hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, hydroxycarbonyl, C<sub>1</sub>-C<sub>6</sub>-alkoxycarbonyl, aminocarbonyl, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, cyano, amino, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylamino,

or

R<sup>6</sup> represents a moiety of the formula



wherein

R<sup>6A</sup> is selected from the group consisting of hydrogen and C<sub>1</sub>-C<sub>4</sub>-alkyl, and

n represents an integer of 1 or 2,

R<sup>7</sup> represents halogen, nitro, cyano, trifluoromethyl, trifluoromethoxy, methyl or ethyl,

and

Y<sup>1</sup>, Y<sup>2</sup>, Y<sup>3</sup>, Y<sup>4</sup> and Y<sup>5</sup> each represent CH.

4. Compounds of general formula (I) according to Claim 1, 2 or 3, wherein

A represents a phenyl or a pyridyl ring,

- 110 -

R<sup>1</sup> and R<sup>3</sup> each represent hydrogen,

R<sup>2</sup> represents fluoro, chloro, bromo, nitro or cyano,


R<sup>4</sup> represents cyano, C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyl or C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl, wherein C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl can be substituted with a radical selected from the group consisting of hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylamino, heteroaryl and heterocyclyl,

R<sup>5</sup> represents methyl,

R<sup>6</sup> represents hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, mono- or di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, C<sub>1</sub>-C<sub>4</sub>-alkylcarbonyl or C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl, wherein C<sub>1</sub>-C<sub>4</sub>-alkyl and C<sub>1</sub>-C<sub>4</sub>-alkoxycarbonyl can be substituted with a radical selected from the group consisting of heteroaryl, hydroxy, C<sub>1</sub>-C<sub>4</sub>-alkoxy, hydroxycarbonyl, aminocarbonyl, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylaminocarbonyl, amino, mono- and di-C<sub>1</sub>-C<sub>4</sub>-alkylamino,

or

R<sup>6</sup> represents a moiety of the formula



wherein

R<sup>6A</sup> is selected from the group consisting of hydrogen and methyl,

- 111 -

$R^7$  represents trifluoromethyl or nitro,

and

5  $Y^1, Y^2, Y^3, Y^4$  and  $Y^5$  each represent CH.

5. Compounds of general formula (I) according to at least one of Claims 1 to 4, wherein A is phenyl or pyridyl.

10 6. Compounds of general formula (I) according to at least one of Claims 1 to 5, wherein  $R^1$  is hydrogen.

7. Compounds of general formula (I) according to at least one of Claims 1 to 6, wherein  $R^2$  is cyano.

15 8. Compounds of general formula (I) according to at least one of Claims 1 to 7, wherein  $R^3$  is hydrogen.

20 9. Compounds of general formula (I) according to at least one of Claims 1 to 8, wherein  $R^4$  is  $C_1$ - $C_4$ -alkoxycarbonyl optionally substituted by hydroxy or wherein  $R^4$  is  $C_1$ - $C_4$ -alkylcarbonyl.


10. Compounds of general formula (I) according to at least one of Claims 1 to 9, wherein  $R^5$  is methyl.

25 11. Compounds of general formula (I) according to at least one of Claims 1 to 10, wherein  $R^6$  is hydrogen.

30 12. Compounds of general formula (I) according to at least one of Claims 1 to 11, wherein  $R^7$  is trifluoromethyl or nitro.

- 112 -

## 13. Compounds of general formula (IA)




wherein

5

Z represents CH or N, and

R<sup>1</sup>, R<sup>3</sup>, R<sup>4</sup> and R<sup>6</sup> have the meaning indicated in Claims 1 to 12.

- 10 14. Process for synthesizing the compounds of general formula (I) or (IA), respectively, as defined in Claims 1 to 13 by condensing compounds of general formula (II)




15

wherein

A, R<sup>1</sup> and R<sup>2</sup> have the meaning indicated in Claims 1 to 13,

with compounds of general formula (III)

20




wherein

R<sup>4</sup> and R<sup>5</sup> have the meaning indicated in Claims 1 to 13,

5

and compounds of general formula (IV)




wherein

10

R<sup>3</sup>, R<sup>7</sup>, and Y<sup>1</sup> to Y<sup>5</sup> have the meaning indicated in Claims 1 to 13,

in the presence of an acid either in a three-component / one-step reaction or sequentially to give compounds of the general formula (IB)

15



wherein

A, R<sup>1</sup> to R<sup>5</sup>, R<sup>7</sup>, and Y<sup>1</sup> to Y<sup>5</sup> have the meaning indicated in Claims 1 to 13,

- 5 optionally followed by reaction of the compounds of general formula (IB) with compounds of the general formula (V)



10 wherein

R<sup>6\*</sup> has the meaning of R<sup>6</sup> as indicated in Claims 1 to 13, but does not represent hydrogen, and

- 15 X represents a leaving group, such as halogen, tosylate, mesylate or sulfate,

in the presence of a base.

- 20 15. The composition containing at least one compound of general formula (I) or (IA) as defined in Claims 1 to 13 and a pharmacologically acceptable diluent.
16. A composition according to Claim 15 for the treatment of acute and chronic inflammatory, ischaemic and/or remodelling processes.
- 25 17. The process for the preparation of compositions according to Claim 15 and 16 characterized in that the compounds of general formula (I) or (IA) as defined in Claims 1 to 13 together with customary auxiliaries are brought into a suitable application form.

30

- 115 -

18. Use of the compounds of general formula (I) or (IA) as defined in Claims 1 to 13 for the preparation of medicaments.
- 5 19. Use according to Claim 18 for the preparation of medicaments for the treatment of acute and chronic inflammatory, ischaemic and/or remodelling processes.
- 10 20. Use according to Claim 19, wherein the process is chronic obstructive pulmonary disease, acute coronary syndrome, acute myocardial infarction or development of heart failure.
- 15 21. Process for controlling chronic obstructive pulmonary disease, acute coronary syndrome, acute myocardial infarction or development of heart failure in humans and animals by administration of a neutrophil elastase inhibitory amount of at least one compound according to any of Claims 1 to 13.



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/09525

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D239/22 C07D401/04 C07D401/12 C07D403/08 C07D403/12  
C07D405/08 C07D413/04 C07D417/04 A61K31/513 A61K31/5377  
A61P9/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	FRECHET J M J ET AL: "A Combinatorial Approach to Recognition of Chirality: Preparation of Highly Enantioselective Aryl-Dihydropyrimidine Selectors for Chiral HPLC" JOURNAL OF COMBINATORIAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, US, vol. 1, no. 1, 1999, pages 105-112, XP002267474 ISSN: 1520-4766 Scheme 1, entry 27 of table 1, page 110, column 2, paragraph 2	1-3,6, 9-11,14, 15
A	EP 0 528 633 A (ICI PLC) 24 February 1993 (1993-02-24) page 2, line 1 - line 18; claims 1,13 -/-	1-21

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the International filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the International filing date but later than the priority date claimed

\*T\* later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*Z\* document member of the same patent family

Date of the actual completion of the International search

21 January 2004

Date of mailing of the International search report

12/02/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Hanisch, I

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/09525

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 01 37837 A (SMITHKLINE BEECHAM CORP ; UNDERWOOD DAVID C (US); ADAMS JERRY L (US) 31 May 2001 (2001-05-31) page 16, line 24 - page 18, line 23; claims 1,12; examples 13,14 -----	1-21
A	US 5 532 366 A (WARNER PETER ET AL) 2 July 1996 (1996-07-02) column 1, line 4 - line 31; claim 1 -----	1-21

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/EP 03/09525

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: —  
because they relate to subject matter not required to be searched by this Authority, namely:  
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claim 21 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

---

Continuation of Box I.1

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/09525

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0528633	A	24-02-1993	AT 197043 T	15-11-2000
			AU 658426 B2	13-04-1995
			AU 2101692 A	18-02-1993
			CA 2076226 A1	16-02-1993
			CZ 9202517 A3	17-02-1993
			DE 69231513 D1	23-11-2000
			DE 69231513 T2	01-03-2001
			EP 0528633 A1	24-02-1993
			FI 923661 A	16-02-1993
			HU 61732 A2	01-03-1993
			IE 922586 A1	24-02-1993
			JP 5286946 A	02-11-1993
			MX 9204712 A1	01-06-1993
			NO 923197 A	16-02-1993
			NZ 243950 A	26-05-1995
			US 5254558 A	19-10-1993
			ZA 9206147 A	28-04-1993
			ZW 13292 A1	05-05-1993
WO 0137837	A	31-05-2001	AU 1626001 A	04-06-2001
			EP 1248624 A1	16-10-2002
			JP 2003517471 T	27-05-2003
			WO 0137837 A1	31-05-2001
US 5532366	A	02-07-1996	AT 171191 T	15-10-1998
			AU 669545 B2	13-06-1996
			AU 4507893 A	31-01-1994
			CA 2139421 A1	20-01-1994
			DE 69321121 D1	22-10-1998
			DE 69321121 T2	04-03-1999
			EP 0649432 A1	26-04-1995
			FI 946202 A	26-01-1995
			WO 9401455 A1	20-01-1994
			HU 68543 A2	28-06-1995
			JP 7508748 T	28-09-1995
			NO 945091 A	16-02-1995